

## **Induction of abortion (2<sup>nd</sup> Trimester)**

### **Referenced to July 2007**

#### **Review Articles**

- Ashok PW, Templeton A, Wagaarachchi PT, Flett GM. Midtrimester medical termination of pregnancy: a review of 1002 consecutive cases. *Contraception* 2004;**69**:51-8.
- Baird DT. Clinical use of mifepristone (RU 486). *Ann.Med.* 1993;**25**:65-9.
- Batioglu S, Tonguc E, Haberal A, Celikkanat H, Bagis T. Midtrimester termination of complicated pregnancy with oral misoprostol. *Adv.Contracept.* 1997;**13**:55-61.
- Bechard dS. [Use of misoprostol in gynecology and obstetrics]. *Gynecol.Obstet.Fertil.* 2002;**30**:317-24.
- Berer M. Medical abortion: Issues of choice and acceptability. *Reproductive Health Matters* 2005;**13**:25-34.
- Berer M. Medical abortion: a fact sheet. *Reprod.Health Matters.* 2005;**13**:20-4.
- Bourguignon A, Briscoe B, Nemzer L. Genetic abortion: Considerations for patient care. *Journal of Perinatal & Neonatal Nursing* 1999;**13**:47-58.
- Dodd JM, Crowther CA. Misoprostol versus cervagem for the induction of labour to terminate pregnancy in the second and third trimester: a systematic review. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2006;**125**:3-8.
- Dommergues M. Termination of pregnancy for fetal neurological abnormalities. *Childs Nervous System* 2003;**19**:600-4.
- Friel L, Rochon M, Gyamfi C, Jacobs A. Safety and efficacy of misoprostol for second-trimester pregnancy termination: effect of prior cesarean section. *American Journal of Obstetrics and Gynecology* 2004;**191**:S85.
- Goldberg AB, Greenberg MB, Darney PD. Misoprostol and pregnancy. *N.Engl.J.Med.* 2001;**344**:38-47.
- Goldberg AB, Greenberg MB, Darney PD. Drug therapy: Misoprostol and pregnancy. *New England Journal of Medicine* 2001;**344**:38-47.
- Lalitkumar S, Bygdeman M, Gemzell-Danielsson K. Mid-trimester induced abortion: a review. *Hum.Reprod.Update.* 2007;**13**:37-52.

Lokugamage AU, Refaey HE, Rodeck CH. Misoprostol and pregnancy: ever-increasing indications of effective usage. *Curr.Opin.Obstet.Gynaecol.* 2003;**15**:513-8.

Moore ML. Misoprostol-is more research needed? *J.Perinat.Educ.* 2002;**11**:43-7.

Ngai SW, Tang OS, Ho PC. Prostaglandins for induction of second-trimester termination and intrauterine death. *Best.Pract.Res.Clin.Obstet.Gynaecol.* 2003;**17**:765-75.

Ramin KD, Ogburn PL, Danilenko DR, Ramsey PS. High-dose oral misoprostol for mid-trimester pregnancy interruption. *Gynecol.Obstet.Invest* 2002;**54**:176-9.

Ramsey PS, Owen J. Midtrimester cervical ripening and labor induction. *Clinical Obstetrics and Gynecology* 2000;**43**:495-512.

Scheepers HC, van Erp EJ, van den Bergh AS. Use of misoprostol in first and second trimester abortion: a review. *Obstet.Gynecol.Surv.* 1999;**54**:592-600.

Schreiber C, Creinin M. Mifepristone in abortion care. *Seminars in Reproductive Medicine* 2005;**23**:82-91.

Shannon C, Brothers LP, Philip NM, Winikoff B. Infection after medical abortion: a review of the literature. *Contraception* 2004;**70**:183-90.

Song J. Use of misoprostol in obstetrics and gynecology. *Obstet.Gynecol.Surv.* 2000;**55**:503-10.

Stubblefield PG, Carr-Ellis S, Borgatta L. Methods for induced abortion. *Obstet.Gynecol.* 2004;**104**:174-85.

Tang OS, Ho PC. Medical abortion in the second trimester. *Best.Pract.Res.Clin.Obstet.Gynaecol.* 2002;**16**:237-46.

Tang OS, Ho PC. Clinical applications of mifepristone. *Gynecological Endocrinology* 2006;**22**:655-9.

Van Look PFA, Cottingham JC. Unsafe abortion: an avoidable tragedy. *Best Practice & Research in Clinical Obstetrics & Gynaecology* 2002;**16**:205-20.

Zust S, Hosli I, Surbek D, Holzgreve W. Side effects of Misoprostol therapy during pregnancy. *Zeitschrift fur Geburtshilfe und Neonatologie* 2001;**205**:43-8.

Zust S, Hosli I, Surbek D, Holzgreve W. [Adverse effects of misoprostol in pregnancy]. *Z.Geburtshilfe Neonatol.* 2001;**205**:43-8.

## Clinical Studies

### Randomised Controlled Trials

Akoury H, Hannah M, Chitayat D, Seaward G, Winsor EJT, Ferris L *et al.* A randomized controlled trial (RCT) to evaluate the use of misoprostol for second trimester pregnancy termination associated with fetal malformation. *American Journal of Obstetrics and Gynecology* 2001;**185**:S112.

Akoury H, Hannah M, Windrim R, Reid DM, Pai A, Thomas M *et al.* A randomized controlled trial (RCT) to evaluate the use of misoprostol for second trimester pregnancy termination associated with fetal malformation. *American Journal of Human Genetics* 2002;**71**:563.

Akoury HA, Hannah ME, Chitayat D, Thomas M, Winsor E, Ferris LE *et al.* Randomized controlled trial of misoprostol for second-trimester pregnancy termination associated with fetal malformation. *Am.J.Obstet.Gynecol.* 2004;**190**:755-62.

Ashok PW, Kidd A, Flett GM, Fitzmaurice A, Graham W, Templeton A. A randomized comparison of medical abortion and surgical vacuum aspiration at 10-13 weeks gestation. *Hum.Reprod.* 2002;**17**:92-8.

Atlas RO, Lemus J, Reed J, Atkins D, Alger LS. Second trimester abortion using prostaglandin E-2 suppositories with or without intracervical *Laminaria japonica*: A randomized study. *Obstetrics and Gynecology* 1998;**92**:398-402.

Bartley J, Baird DT. A randomised study of misoprostol and gemeprost in combination with mifepristone for induction of abortion in the second trimester of pregnancy. *BJOG.* 2002;**109**:1290-4.

Bebbington MW, Kent N, Lim K, Gagnon A, Delisle MF, Tessier F *et al.* A randomized controlled trial comparing two protocols for the use of misoprostol in midtrimester pregnancy termination. *Am.J.Obstet.Gynecol.* 2002;**187**:853-7.

Bhattacharyya SK, Mukherji J, Kamilya GS, Ray S, Hazra A. Two regimens of vaginal misoprostol in second trimester termination of pregnancy: a prospective randomised trial. *Acta Obstet.Gynecol.Scand.* 2006;**85**:1458-62.

Borgatta L, Chen AY, Vragovic O, Stubblefield PG, Magloire CA. A randomized clinical trial of the addition of laminaria to misoprostol and hypertonic saline for second-trimester induction abortion. *Contraception* 2005;**72**:358-61.

Bugalho A, Faundes A, Jamisse L, Usfa M, Maria E, Bique C. Evaluation of the effectiveness of vaginal misoprostol to induce first trimester abortion. *Contraception* 1996;**53**:244-6.

Butt K, Delaney T, Hutchens D, Crane J, Young D. Randomized comparison of oral and vaginal misoprostol for midtrimester pregnancy termination for fetal demise and major congenital anomalies. *American Journal of Obstetrics and Gynecology* 2001;**185**:S205.

Caliskan E, Dilbaz S, Doger E, Ozeren S, Dilbaz B. Randomized comparison of 3 misoprostol protocols for abortion induction at 13-20 weeks of gestation. *J.Reprod.Med.* 2005;**50**:173-80.

Cheng L. [Termination of 10-16 weeks's gestation with mifepristone plus misoprostol: a multicentre randomized clinical trial]. *Zhonghua Fu Chan Ke.Za Zhi.* 1999;**34**:268-71.

Dickinson, Evans. The optimization of intravaginal misoprostol dosing schedules in second-trimester pregnancy termination (vol 186, pg 470, 2002). *American Journal of Obstetrics and Gynecology* 2005;**193**:597.

Dickinson JE, Godfrey M, Evans SF. Efficacy of intravaginal misoprostol in second-trimester pregnancy termination: a randomized controlled trial. *J.Matern.Fetal Med.* 1998;**7**:115-9.

Dickinson JE, Evans SF. The optimization of intravaginal misoprostol dosing schedules in second-trimester pregnancy termination. *Am.J.Obstet.Gynecol.* 2002;**186**:470-4.

Dickinson JE, Evans SF. A comparison of oral misoprostol with vaginal misoprostol administration in second-trimester pregnancy termination for fetal abnormality. *Obstet.Gynecol.* 2003;**101**:1294-9.

Donovan P. Misoprostol May Induce 2Nd-Trimester Abortion with Fewer Side-Effects Than Other Prostaglandins. *Family Planning Perspectives* 1995;**27**:43-4.

El Refaey H, Hinshaw K, Templeton A. The abortifacient effect of misoprostol in the second trimester. A randomized comparison with gemeprost in patients pre-treated with mifepristone (RU486). *Hum.Reprod.* 1993;**8**:1744-6.

El Refaey H, Templeton A. Induction of abortion in the second trimester by a combination of misoprostol and mifepristone: a randomized comparison between two misoprostol regimens. *Hum.Reprod.* 1995;**10**:475-8.

Feldman DM, Borgida AF, Rodis JF, Leo MV, Campbell WA. A randomized comparison of two regimens of misoprostol for second-trimester pregnancy termination. *Am.J.Obstet.Gynecol.* 2003;**189**:710-3.

Ghorab MN, El Helw BA. Second-trimester termination of pregnancy by extra-amniotic prostaglandin F2alpha or endocervical misoprostol. A comparative study. *Acta Obstet.Gynecol.Scand.* 1998;**77**:429-32.

- Gilbert A, Reid R. A randomised trial of oral versus vaginal administration of misoprostol for the purpose of mid-trimester termination of pregnancy. *Aust.N.Z.J.Obstet.Gynaecol.* 2001;**41**:407-10.
- Gonzalez JA, Carlan SJ, Alverson MW. Outpatient second trimester pregnancy termination. *Contraception* 2001;**63**:89-93.
- Grimes DA, Smith MS, Witham AD. Mifepristone and misoprostol versus dilation and evacuation for midtrimester abortion: a pilot randomised controlled trial. *BJOG.* 2004;**111**:148-53.
- Guix C, Palacio M, Figueras F, Bennasar M, Zamora L, Coll O *et al.* Efficacy of two regimens of misoprostol for early second-trimester pregnancy termination. *Fetal Diagn.Ther.* 2005;**20**:544-8.
- Hamoda H, Ashok PW, Flett GM, Templeton A. A randomised controlled trial of mifepristone in combination with misoprostol administered sublingually or vaginally for medical abortion up to 13 weeks of gestation. *BJOG.* 2005;**112**:1102-8.
- Hamoda H, Ashok PW, Flett GM, Templeton A. A randomized trial of mifepristone in combination with misoprostol administered sublingually or vaginally for medical abortion at 13-20 weeks gestation. *Hum.Reprod.* 2005;**20**:2348-54.
- Heimstad R, Backe B. [Mifepristone for induction of second trimester abortion]. *Tidsskr.Nor Laegeforen.* 2003;**123**:2412-3.
- Herabutya Y, Chanrachakul B, Punyavachira P. Second trimester pregnancy termination: a comparison of 600 and 800 micrograms of intravaginal misoprostol. *J.Obstet.Gynaecol.Res.* 2001;**27**:125-8.
- Herabutya Y, Chanrachakul B, Punyavachira P. A randomised controlled trial of 6 and 12 hourly administration of vaginal misoprostol for second trimester pregnancy termination. *BJOG.* 2005;**112**:1297-301.
- Hidar S, Fekih M, Chaieb A, Bibi M, Mellouli R, Khairi H. [Oxytocin and misoprostol administered intravaginally for termination of pregnancy at 13 to 29 weeks of amenorrhea. A prospective randomized trial]. *J.Gynecol.Obstet.Biol.Reprod.(Paris)* 2001;**30**:439-43.
- Hidar S, Bouddebous M, Chaieb A, Jerbi M, Bibi M, Khairi H. Randomized controlled trial of vaginal misoprostol versus vaginal misoprostol and isosorbide dinitrate for termination of pregnancy at 13-29 weeks. *Arch.Gynecol.Obstet.* 2005;**273**:157-60.
- Ho PC, Chan YF, Lau W. Misoprostol is as effective as gemeprost in termination of second trimester pregnancy when combined with mifepristone: a randomised comparative trial. *Contraception* 1996;**53**:281-3.

- Ho PC, Ngai SW, Liu KL, Wong GC, Lee SW. Vaginal misoprostol compared with oral misoprostol in termination of second-trimester pregnancy. *Obstet.Gynecol.* 1997;**90**:735-8.
- Jain JK, Mishell DR, Jr. A comparison of intravaginal misoprostol with prostaglandin E2 for termination of second-trimester pregnancy. *N.Engl.J.Med.* 1994;**331**:290-3.
- Jain JK, Mishell DR, Jr. A comparison of misoprostol with and without laminaria tents for induction of second-trimester abortion. *Am.J.Obstet.Gynecol.* 1996;**175**:173-7.
- Jain JK, Kuo J, Mishell DR, Jr. A comparison of two dosing regimens of intravaginal misoprostol for second-trimester pregnancy termination. *Obstet.Gynecol.* 1999;**93**:571-5.
- Kelekci S, Erdemoglu E, Inan I. Randomized study on the effect of adding oxytocin to ethacridine lactate or misoprostol for second-trimester termination of pregnancy. *Acta Obstet.Gynecol.Scand.* 2006;**85**:825-9.
- Khazardoost S, Hantoushzadeh S, Madani MM. A randomised trial of two regimens of vaginal misoprostol to manage termination of pregnancy of up to 16 weeks. *Aust.N.Z.J.Obstet.Gynaecol.* 2007;**47**:226-9.
- Kmak D, Refuerzo J, Johnson A, Blackwell S, McNeeley G, Qureshi F *et al.* Comparison of oral, intravaginal and combination misoprostol therapy for second trimester inductions. *American Journal of Obstetrics and Gynecology* 2001;**185**:S205.
- le Roux PA, Tregoning SK, Zinn PM, van der Spuy ZM. Inhibition of progesterone secretion with trilostane for mid-trimester termination of pregnancy: randomized controlled trials. *Hum.Reprod.* 2002;**17**:1483-9.
- le Roux PA, van der Spuy ZM. Labor induction abortion utilizing trilostane, a 3beta-hydroxysteroid dehydrogenase inhibitor. *Contraception* 2005;**71**:343-7.
- Leader J, Bujnovsky M, Carlan SJ, Triana T, Richichi K. Effect of oral misoprostol after second-trimester delivery: a randomized, blinded study. *Obstet.Gynecol.* 2002;**100**:689-94.
- Lemyre M, Leboeuf M, Turcot-Lemay L, Thomas N. A randomized study comparing two regimens of intravaginal misoprostol for midtrimester medical pregnancy termination. *American Journal of Obstetrics and Gynecology* 2006;**195**:S105.
- Li CF, Chan CW, Ho PC. A study of the efficacy of cervical ripening with nitric oxide donor versus placebo for cervical priming before second-trimester termination of pregnancy. *Contraception* 2003;**68**:269-72.

Makhlouf AM, Al Hussaini TK, Habib DM, Makarem MH. Second-trimester pregnancy termination: comparison of three different methods. *J.Obstet.Gynaecol.* 2003;**23**:407-11.

Marquette GP, Skoll MA, Dontigny L. A randomized trial comparing oral misoprostol with intra-amniotic prostaglandin F2alpha for second trimester terminations. *J.Obstet.Gynaecol.Can.* 2005;**27**:1013-8.

Munthali J, Moodley J. The use of misoprostol for mid-trimester therapeutic termination of pregnancy. *Trop.Doct.* 2001;**31**:157-61.

Ngai SW, Tang OS, Ho PC. Randomized comparison of vaginal (200 microg every 3 h) and oral (400 microg every 3 h) misoprostol when combined with mifepristone in termination of second trimester pregnancy. *Hum.Reprod.* 2000;**15**:2205-8.

Nigam A, Singh VK, Prakash A. Vaginal vs. oral misoprostol for mid-trimester abortion. *Int.J.Gynaecol.Obstet.* 2006;**92**:270-1.

Niromanesh S, Hashemi-Fesharaki M, Mosavi-Jarrahi A. Second trimester abortion using intravaginal misoprostol. *Int.J.Gynaecol.Obstet.* 2005;**89**:276-7.

Nor Azlin MI, Abdullah HS, Zainul Rashid MR, Jamil MA. Misoprostol (alone) in second trimester terminations of pregnancy: as effective as Gemeprost? *J.Obstet.Gynaecol.* 2006;**26**:546-9.

Nuthalapaty FS, Ramsey PS, Biggio JR, Owen J. High-dose vaginal misoprostol versus concentrated oxytocin plus low-dose vaginal misoprostol for midtrimester labor induction: a randomized trial. *Am.J.Obstet.Gynecol.* 2005;**193**:1065-70.

Nuutila M, Toivonen J, Ylikorkala O, Halmesmaki E. A comparison between two doses of intravaginal misoprostol and gemeprost for induction of second-trimester abortion. *Obstet.Gynecol.* 1997;**90**:896-900.

Owen J, Hauth JC. Vaginal misoprostol vs. concentrated oxytocin plus low-dose prostaglandin E2 for second trimester pregnancy termination. *J.Matern.Fetal Med.* 1999;**8**:48-50.

Paz B, Ohel G, Tal T, Degani S, Sabo E, Levitan Z. Second trimester abortion by laminaria followed by vaginal misoprostol or intrauterine prostaglandin F2alpha: a randomized trial. *Contraception* 2002;**65**:411-3.

Perry KG, Jr., Rinehart BK, Terrone DA, Martin RW, May WL, Roberts WE. Second-trimester uterine evacuation: A comparison of intra-amniotic (15S)-15-methyl-prostaglandin F2alpha and intravaginal misoprostol. *Am.J.Obstet.Gynecol.* 1999;**181**:1057-61.

- Ramsey PS, Savage K, Lincoln T, Owen J. Vaginal misoprostol versus concentrated oxytocin and vaginal PGE2 for second-trimester labor induction. *Obstet.Gynecol.* 2004;**104**:138-45.
- Roy G, Ferreira E, Hudon L, Marquette G. The efficacy of oral versus vaginal misoprostol for second-trimester termination of pregnancy: A double-blind, randomized, placebo-controlled trial. *American Journal of Obstetrics and Gynecology* 2003;**189**:S70.
- Ruangchainikhom W, Phongphissanou E, Bhekasuta J, Sarapak S. Effectiveness of 400 or 600 micrograms of vaginal misoprostol for terminations of early pregnancies. *J.Med.Assoc.Thai.* 2006;**89**:928-33.
- Saha S, Bal R, Ghosh S, Krishnamurthy P. Medical abortion in late second trimester-- a comparative study with misoprostol through vaginal versus oral followed by vaginal route. *J.Indian Med.Assoc.* 2006;**104**:81-2, 84.
- Su LL, Biswas A, Choolani M, Kalaichelvan V, Singh K. A prospective, randomized comparison of vaginal misoprostol versus intra-amniotic prostaglandins for midtrimester termination of pregnancy. *Am.J.Obstet.Gynecol.* 2005;**193**:1410-4.
- Tang OS, Lau WN, Chan CC, Ho PC. A prospective randomised comparison of sublingual and vaginal misoprostol in second trimester termination of pregnancy. *BJOG.* 2004;**111**:1001-5.
- Tang OS, Chan CC, Kan AS, Ho PC. A prospective randomized comparison of sublingual and oral misoprostol when combined with mifepristone for medical abortion at 12-20 weeks gestation. *Hum.Reprod.* 2005;**20**:3062-6.
- Webster D, Penney GC, Templeton A. A comparison of 600 and 200 mg mifepristone prior to second trimester abortion with the prostaglandin misoprostol. *Br.J.Obstet.Gynaecol.* 1996;**103**:706-9.
- Wong KS, Ngai CS, Chan KS, Tang LC, Ho PC. Termination of second trimester pregnancy with gemeprost and misoprostol: a randomized double-blind placebo-controlled trial. *Contraception* 1996;**54**:23-5.
- Wong KS, Ngai CS, Wong AY, Tang LC, Ho PC. Vaginal misoprostol compared with vaginal gemeprost in termination of second trimester pregnancy. A randomized trial. *Contraception* 1998;**58**:207-10.
- Wong KS, Ngai CS, Yeo EL, Tang LC, Ho PC. A comparison of two regimens of intravaginal misoprostol for termination of second trimester pregnancy: a randomized comparative trial. *Hum.Reprod.* 2000;**15**:709-12.
- Yapar EG, Senoz S, Urkudur M, Batioglu S, Gokmen O. Second trimester pregnancy termination including fetal death: comparison of five different methods. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 1996;**69**:97-102.

Yilmaz B, Kelekci S, Ertas IE, Kahyaoglu S, Ozel M, Sut N *et al.* Misoprostol moistened with acetic acid or saline for second trimester pregnancy termination: a randomized prospective double-blind trial. *Hum.Reprod.* 2005;**20**:3067-71.

Yilmaz B, Kelekci S, Ertas IE, Ozel M, Sut N, Mollamahmutoglu L *et al.* Randomized comparison of second trimester pregnancy termination utilizing saline moistened or dry misoprostol. *Arch.Gynecol.Obstet.* 2007.

## Observational Studies

Al Hussaini TK. Uterine rupture in second trimester abortion in a grand multiparous woman. A complication of misoprostol and oxytocin. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2001;**96**:218-9.

Al Taani MI. Termination of second trimester, complicated gestation. *East Mediterr.Health J.* 2005;**11**:657-62.

Ashok PW, Flett GM, Templeton A. Termination of pregnancy at 9-13 weeks' amenorrhoea with mifepristone and misoprostol. *Lancet* 1998;**352**:542-3.

Ashok PW, Templeton A. Nonsurgical mid-trimester termination of pregnancy: a review of 500 consecutive cases. *Br.J.Obstet.Gynaecol.* 1999;**106**:706-10.

Autry AM, Hayes EC, Jacobson GF, Kirby RS. A comparison of medical induction and dilation and evacuation for second-trimester abortion. *Am.J.Obstet.Gynecol.* 2002;**187**:393-7.

Bagga R, Chaudhary N, Kalra J. Rupture in an unscarred uterus during second trimester pregnancy termination with mifepristone and misoprostol. *Int.J.Gynaecol.Obstet.* 2004;**87**:42-3.

Bartlett LA, Berg CJ, Shulman HB, Zane SB, Green CA, Whitehead S *et al.* Risk factors for legal induced abortion-related mortality in the United States. *Obstetrics and Gynecology* 2004;**103**:729-37.

Basu R, Gundlach T, Tasker M. Mifepristone and misoprostol for medical termination of pregnancy: the effectiveness of a flexible regimen. *J.Fam.Plann.Reprod.Health Care* 2003;**29**:139-41.

Berghahn L, Christensen D, Droste S. Uterine rupture during second-trimester abortion associated with misoprostol. *Obstet.Gynecol.* 2001;**98**:976-7.

Bhattacharjee N, Ganguly RP, Saha SP. Misoprostol for termination of mid-trimester post-Caesarean pregnancy. *Aust.N.Z.J.Obstet.Gynaecol.* 2007;**47**:23-5.

Bracken H, Ngoc NT, Schaff E, Coyaji K, Ambardekar S, Westheimer E *et al.* Mifepristone followed in 24 hours to 48 hours by misoprostol for late first-trimester abortion. *Obstet.Gynecol.* 2007;**109**:895-901.

Bugalho A, Bique C, Almeida L, Bergstrom S. Pregnancy interruption by vaginal misoprostol. *Gynecol.Obstet.Invest* 1993;**36**:226-9.

Bugalho A, Bique C, Almeida L, Faundes A. The effectiveness of intravaginal misoprostol (Cytotec) in inducing abortion after eleven weeks of pregnancy. *Stud.Fam.Plann.* 1993;**24**:319-23.

Bugalho A, Bique C, Pereira C, Granja AC, Bergstrom S. Uterine evacuation by vaginal misoprostol after second trimester pregnancy interruption. *Acta Obstet.Gynecol.Scand.* 1996;**75**:270-3.

Carbonell JL, Varela L, Velazco A, Fernandez C. The use of misoprostol for termination of early pregnancy. *Contraception* 1997;**55**:165-8.

Carbonell JL, Valera L, Velazco A, Tanda R, Sanchez C. Vaginal misoprostol for early second-trimester abortion. *Eur.J.Contracept.Reprod.Health Care* 1998;**3**:93-8.

Carbonell JL, Varela L, Velazco A, Tanda R, Sanchez C. Vaginal misoprostol for abortion at 10-13 weeks' gestation. *Eur.J.Contracept.Reprod.Health Care* 1999;**4**:35-40.

Carbonell JL, Velazco A, Varela L, Tanda R, Sanchez C, Barambio S *et al.* Misoprostol for abortion at 9-12 weeks' gestation in adolescents. *Eur.J.Contracept.Reprod.Health Care* 2001;**6**:39-45.

Carbonell JL, Rodriguez J, Delgado E, Sanchez C, Vargas F, Valera L *et al.* Vaginal misoprostol 800 microg every 12 h for second-trimester abortion. *Contraception* 2004;**70**:55-60.

Chen M, Shih JC, Chiu WT, Hsieh FJ. Separation of cesarean scar during second-trimester intravaginal misoprostol abortion. *Obstet.Gynecol.* 1999;**94**:840.

Choudhary N, Saha SC, Gopalan S. Abortion procedures in a tertiary care institution in India. *Int.J.Gynaecol.Obstet.* 2005;**91**:81-6.

Clark S, Blum J, Blanchard K, Galvao L, Fletcher H, Winikoff B. Misoprostol use in obstetrics and gynecology in Brazil, Jamaica, and the United States. *Int.J.Gynaecol.Obstet.* 2002;**76**:65-74.

Costa SH, Vessey MP. Misoprostol and illegal abortion in Rio de Janeiro, Brazil. *Lancet* 1993;**341**:1258-61.

Cowett AA, Golub RM, Grobman WA. Cost-effectiveness of dilation and evacuation versus the induction of labor for second-trimester pregnancy termination. *Am.J.Obstet.Gynecol.* 2006;**194**:768-73.

Daponte A, Nzewenga G, Dimopoulos KD, Guidozi F. The use of vaginal misoprostol for second-trimester pregnancy termination in women with previous single cesarean section. *Contraception* 2006;**74**:324-7.

Daskalakis G, Papantoniou N, Mesogitis S, Papageorgiou J, Antsaklis A. Sonographic findings and surgical management of a uterine rupture associated with the use of misoprostol during second-trimester abortion. *J.Ultrasound Med.* 2005;**24**:1565-8.

Daskalakis GJ, Mesogitis SA, Papantoniou NE, Mouloupoulos GG, Papapanagiotou AA, Antsaklis AJ. Misoprostol for second trimester pregnancy termination in women with prior caesarean section. *BJOG.* 2005;**112**:97-9.

de Boer MA, van Gemund N, Scherjon SA, Kanhai HHH. Low dose sulprostone for termination of second and third trimester pregnancies. *European Journal of Obstetrics Gynecology and Reproductive Biology* 2001;**99**:244-8.

De Heus R, Graziosi GC, Christiaens GC, Bruinse HW, Mol BW. Medical management for termination of second and third trimester pregnancies: a comparison of strategies. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2004;**116**:16-21.

Dickinson JE. Late pregnancy termination within a legislated medical environment. *Aust.N.Z.J.Obstet.Gynaecol.* 2004;**44**:337-41.

Dickinson JE. Misoprostol for second-trimester pregnancy termination in women with a prior cesarean delivery. *Obstet.Gynecol.* 2005;**105**:352-6.

Dilbaz S, Caliskan E, Dilbaz B, Kahraman BG. Frequent low-dose misoprostol for termination of second-trimester pregnancy. *Eur.J.Contracept.Reprod.Health Care* 2004;**9**:11-5.

Dodd J, O'Brien L, Coffey J. Misoprostol for second and third trimester termination of pregnancy: a review of practice at the Women's and Children's Hospital, Adelaide, Australia. *Aust.N.Z.J.Obstet.Gynaecol.* 2005;**45**:25-9.

Edwards RK, Sims SM. Outcomes of second-trimester pregnancy terminations with misoprostol: comparing 2 regimens. *Am.J.Obstet.Gynecol.* 2005;**193**:544-8.

El Matary A, Navaratnarajah R, Economides DL. Ultrasound diagnosis of uterine dehiscence following mifepristone/misoprostol regime in early second trimester termination. *J.Obstet.Gynaecol.* 2006;**26**:578-80.

Elsheikh A, Antsaklis A, Mesogitis S, Papantoniou N, Rodolakis A, Vogas E *et al.* Use of misoprostol for the termination of second trimester pregnancies. *Arch.Gynecol.Obstet.* 2001;**265**:204-6.

- Garbin O, Vayssiere C, Bettahar-Lebugle K, Nisand I. Consistency of medical abortion efficacy from 5 through 14 weeks' gestation. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2006;**129**:36-40.
- Goh SE, Thong KJ. Induction of second trimester abortion (12-20 weeks) with mifepristone and misoprostol: a review of 386 consecutive cases. *Contraception* 2006;**73**:516-9.
- Gouk EV, Lincoln K, Khair A, Haslock J, Knight J, Cruickshank DJ. Medical termination of pregnancy at 63 to 83 days gestation. *Br.J.Obstet.Gynaecol.* 1999;**106**:535-9.
- Hamoda H, Ashok PW, Flett GM, Templeton A. Medical abortion at 64 to 91 days of gestation: a review of 483 consecutive cases. *Am.J.Obstet.Gynecol.* 2003;**188**:1315-9.
- Hamoda H, Ashok PW, Flett GM, Templeton A. Analgesia requirements and predictors of analgesia use for women undergoing medical abortion up to 22 weeks of gestation. *BJOG.* 2004;**111**:996-1000.
- Hamoda H, Ashok PW, Flett GM, Templeton A. Medical abortion at 9-13 weeks' gestation: a review of 1076 consecutive cases. *Contraception* 2005;**71**:327-32.
- Heikinheimo O, Suhonen S, Haukkamaa M. One- and 2-day mifepristone-misoprostol intervals are both effective in medical termination of second-trimester pregnancy. *Reprod.Biomed.Online.* 2004;**8**:236-9.
- Herabutya Y, Prasertsawat P. Second trimester abortion using intravaginal misoprostol. *Int.J.Gynaecol.Obstet.* 1998;**60**:161-5.
- Herabutya Y, Chanrachakul B, Punyavachira P. Vaginal misoprostol in termination of second trimester pregnancy. *J.Obstet.Gynaecol.Res.* 2000;**26**:121-5.
- Herabutya Y, Chanarachakul B, Punyavachira P. Induction of labor with vaginal misoprostol for second trimester termination of pregnancy in the scarred uterus. *Int.J.Gynaecol.Obstet.* 2003;**83**:293-7.
- Hern WM. Misoprostol as an adjunctive medication in late surgical abortion. *Int.J.Gynaecol.Obstet.* 2005;**88**:327-8.
- Hinshaw K, El Refaey H, Rispin R, Templeton A. Mid-trimester termination for fetal abnormality: advantages of a new regimen using mifepristone and misoprostol. *Br.J.Obstet.Gynaecol.* 1995;**102**:559-60.
- Hoffer MC, Charlier C, Giacalone PL, Zimbris L, Astruc M, Boulot P. [Evaluation of combination RU 486-laminaria tents-misoprostol-peridural anesthesia in second and third trimester induced abortions]. *J.Gynecol.Obstet.Biol.Reprod.(Paris)* 1998;**27**:83-6.

- Hsieh TY, Yu CH, Kuo PL, Chang FM. Prenatal diagnosis of alobar holoprosencephaly with cystic hygroma. *Taiwan.J.Obstet.Gynecol.* 2006;**45**:146-9.
- Hunter A, Porter H, Kyle P. Not assuming the obvious: failed surgical termination of pregnancy and multiple fetal abnormalities. *BJOG.* 2002;**109**:1069-71.
- Jannet D, Aflak N, Abankwa A, Carbonne B, Marpeau L, Milliez J. Termination of 2nd and 3rd trimester pregnancies with mifepristone and misoprostol. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 1996;**70**:159-63.
- Jannet D, Aflak N, Chemla JP, Marpeau L, Milliez J. [Mifepristone and misoprostol in therapeutic abortions]. *J.Gynecol.Obstet.Biol.Reprod.(Paris)* 1996;**25**:277-82.
- Khomasuridze AG, Tsertsvadze GL, Parculi MG, Parunashvili NR, Sulamanidze S. [Antiprogesterone for medical abortion]. *Georgian.Med.News* 2006;24-8.
- Langer BR, Peter C, Firtion C, David E, Haberstick R. Second and third medical termination of pregnancy with misoprostol without mifepristone. *Fetal Diagn.Ther.* 2004;**19**:266-70.
- le Roux PA, Pahal GS, Hoffman L, Nooh R, El Refaey H, Rodeck CH. Second trimester termination of pregnancy for fetal anomaly or death: comparing mifepristone/misoprostol to gemeprost. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2001;**95**:52-4.
- Letourneur B, Parant O, Tofani V, Berrebi A. [Uterine rupture on unscarred uterus following labor induction for 2(nd) trimester termination of pregnancy with oral misoprostol: conservative management]. *J.Gynecol.Obstet.Biol.Reprod.(Paris)* 2002;**31**:371-3.
- Li YT, Yin CS. Delivery of retained placenta by misoprostol in second trimester abortion. *Int.J.Gynaecol.Obstet.* 2001;**74**:215-6.
- Liaquat NF, Javed I, Shuja S, Shoaib T, Bano K, Waheed S *et al.* Therapeutic termination of second trimester pregnancies with low dose misoprostol. *J.Coll.Physicians Surg.Pak.* 2006;**16**:464-7.
- Lim JM, Soh EB, Raman S. Intravaginal misoprostol for termination of midtrimester pregnancy. *Aust.N.Z.J.Obstet.Gynaecol.* 1995;**35**:54-5.
- Makrydimas G, Papanikolaou E, Paraskevaidis E, Paschopoulos M, Lolis D. Upper limb abnormalities as an isolated ultrasonographic finding in early detection of trisomy 18. A case report. *Fetal Diagn.Ther.* 2003;**18**:401-3.
- Mazouni C, Provensal M, Porcu G, Guidicelli B, Heckenroth H, Gamberre M *et al.* Termination of pregnancy in patients with previous cesarean section. *Contraception* 2006;**73**:244-8.

- Mendilcioglu I, Simsek M, Seker PE, Erbay O, Zorlu CG, Trak B. Misoprostol in second and early third trimester for termination of pregnancies with fetal anomalies. *Int.J.Gynaecol.Obstet.* 2002;**79**:131-5.
- Nayki U, Taner CE, Mizrak T, Nayki C, Derin G. Uterine rupture during second trimester abortion with misoprostol. *Fetal Diagn.Ther.* 2005;**20**:469-71.
- Naz S, Sultana N. Role of misoprostol for therapeutic termination of pregnancy from 10 -28 weeks of gestation. *J.Pak.Med.Assoc.* 2007;**57**:129-32.
- Ojidu JJ, Sabharwal SD. Setting up a one-stop mifepristone-misoprostol medical termination of pregnancy service for all gestations from 5 to 23 weeks--a review of 482 cases. *J.Obstet.Gynaecol.* 2001;**21**:386-8.
- Oteri O, Hopkins R. Second trimester therapeutic abortion using mifepristone and oral misoprostol in a woman with two previous caesarean sections and a cone biopsy. *J.Matern.Fetal Med.* 1999;**8**:300-1.
- Pappa KI, Anagnou NP, Elsheikh A, Bikouvarakis SS, Konstantinidou A, Salamalekis E. Congenital bronchopulmonary sequestration presenting as a thoracic tumor: a case report. *Eur.J.Gynaecol.Oncol.* 2004;**25**:749-51.
- Petri AL. [Uterine rupture during second trimester abortion induced with misoprostol]. *Ugeskr.Laeger* 2003;**165**:2894-5.
- Phillips K, Berry C, Mathers AM. Uterine rupture during second trimester termination of pregnancy using mifepristone and a prostaglandin. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 1996;**65**:175-6.
- Pongsatha S, Tongsong T. Second trimester pregnancy termination with 800 mcg vaginal misoprostol. *J.Med.Assoc.Thai.* 2001;**84**:859-63.
- Pongsatha S, Tongsong T, Suwannawut O. Therapeutic termination of second trimester pregnancy with vaginal misoprostol. *J.Med.Assoc.Thai.* 2001;**84**:515-9.
- Pongsatha S, Morakot N, Tongsong T. Demographic characteristics of women with self use of misoprostol for pregnancy interruption attending Maharaj Nakorn Chiang Mai Hospital. *J.Med.Assoc.Thai.* 2002;**85**:1074-80.
- Pongsatha S, Tongsong T. Misoprostol for second trimester termination of pregnancies with prior low transverse cesarean section. *Int.J.Gynaecol.Obstet.* 2003;**80**:61-2.
- Pongsatha S, Tongsong T. Second-trimester pregnancy interruption with vaginal misoprostol in women with previous cesarean section. *J.Med.Assoc.Thai.* 2006;**89**:1097-100.

Prachasilpchai N, Russameecharoen K, Borriboonhirunsarn D. Success rate of second-trimester termination of pregnancy using misoprostol. *J.Med.Assoc.Thai.* 2006;**89**:1115-9.

Ramsey PS, Hogg BB, Savage KG, Winkler DD, Owen J. Cardiovascular effects of intravaginal misoprostol in the mid trimester of pregnancy. *Am.J.Obstet.Gynecol.* 2000;**183**:1100-2.

Rose SB, Shand C, Simmons A. Mifepristone- and misoprostol-induced mid-trimester termination of pregnancy: a review of 272 cases. *Aust.N.Z.J.Obstet.Gynaecol.* 2006;**46**:479-85.

Rouzi AA. Second-trimester pregnancy termination with misoprostol in women with previous cesarean sections. *Int.J.Gynaecol.Obstet.* 2003;**80**:317-8.

Ruano R, Dumez Y, Cabrol D, Dommergues M. Second- and third-trimester therapeutic terminations of pregnancy in cases with complete placenta previa--does feticide decrease postdelivery maternal hemorrhage? *Fetal Diagn.Ther.* 2004;**19**:475-8.

Shammas AG, Momani MD. Misoprostol for termination of second trimester pregnancy in a scarred uterus. *Saudi.Med.J.* 2006;**27**:1173-6.

Simcox R, Ash AK, Shennan AH. Fetal extrusion into the lower segment: a cause of failed misoprostol induced midtrimester termination. *J.Obstet.Gynaecol.* 2006;**26**:174.

Sirimai K, Kiriwat O, Neungton S, Suvanichchati S. Misoprostal use for therapeutic abortion in Siriraj Hospital: the year 2000. *J.Med.Assoc.Thai.* 2002;**85**:416-23.

Srisomboon J, Tongsong T, Pongpisuttinun S. Termination of second-trimester pregnancy with intracervicovaginal misoprostol. *J.Med.Assoc.Thai.* 1997;**80**:242-6.

Srisomboon J, Pongpisuttinun S. Efficacy of intracervicovaginal misoprostol in second-trimester pregnancy termination: a comparison between live and dead fetuses. *J.Obstet.Gynaecol.Res.* 1998;**24**:1-5.

Stoutjesdijk JA, Alsmas J, Wildschut HI. [Pregnancy termination in the second trimester with vaginally administered dinoprostone followed by intravenous sulprostone, for the indication 'foetal congenital defects'; results of a retrospective study]. *Ned.Tijdschr.Geneeskd.* 2004;**148**:1293-6.

Tang OS, Ho PC. Pilot study on the use of sublingual misoprostol for medical abortion. *Contraception* 2001;**64**:315-7.

Tarim E, Kilicdag E, Bagis T, Ilgin A, Yanik F. Second-trimester pregnancy termination with oral misoprostol in women who have had one cesarean section. *Int.J.Gynaecol.Obstet.* 2005;**90**:84-5.

Todd CS, Soler M, Castleman L, Rogers MK, Blumenthal PD. Buccal misoprostol as cervical preparation for second trimester pregnancy termination. *Contraception* 2002;**65**:415-8.

Verhulsdonk MT, Papatsonis DN, Heydanus R. [Results of pregnancy termination for foetal congenital or chromosomal disorders or intrauterine foetal death by oral mifepristone and/or intravaginal misoprostol; results of a retrospective study]. *Ned.Tijdschr.Geneeskd.* 2007;**151**:815-20.

Vimala N, Dadhwal V, Mittal S. Sublingual misoprostol for second-trimester abortion. *Int.J.Gynaecol.Obstet.* 2004;**84**:89-90.

Weeks AD, Stewart P. The Use of Mifepristone in Combination with Misoprostol for 2Nd Trimester Termination of Pregnancy. *British Journal of Family Planning* 1995;**21**:43-4.

Whitecar PW, Zweifel A, Moise KJ, Jr. Failed second-trimester misoprostol termination responding to vaginal instillation of citric acid. *Obstet.Gynecol.* 1999;**94**:839.

Willems FE. [Pregnancy termination in the second trimester with vaginally administered dinoprostone followed by intravenous sulprostone, for the indication 'foetal congenital defects'; results of a retrospective study]. *Ned.Tijdschr.Geneeskd.* 2004;**148**:1656-7.

Wright-Francis DL, Raynor BD, Webb GW. Misoprostol in second trimester termination of pregnancy. *Prim.Care Update.Ob Gyns.* 1998;**5**:176.

Yacoub A, Martel MJ. Pregnancy with primary dilated cardiomyopathy. *Obstet.Gynecol.* 2002;**99**:928-30.

Zeteroglu S, Kotan C, Ozen S, Goktolga U. Mucinous appendicular cystadenocarcinoma during pregnancy. A case report. *J.Reprod.Med.* 2003;**48**:831-3.

## **Cost-effectiveness**

Cowett AA, Golub RM, Grobman WA. Cost-effectiveness of dilation and evacuation versus the induction of labor for second-trimester pregnancy termination. *Am.J.Obstet.Gynecol.* 2006;**194**:768-73.

## Pharmacology

Goldberg AB, Greenberg MB, Darney PD. Misoprostol and pregnancy. *N.Engl.J.Med.* 2001;**344**:38-47.

Zust S, Hosli I, Surbek D, Holzgreve W. Side effects of Misoprostol therapy during pregnancy. *Zeitschrift fur Geburtshilfe und Neonatologie* 2001;**205**:43-8.

Zust S, Hosli I, Surbek D, Holzgreve W. [Adverse effects of misoprostol in pregnancy]. *Z.Geburtshilfe Neonatol.* 2001;**205**:43-8.

## Politics/ Licensing/ Ethics

[Anon]. USFDA, pharmacia new ruling on misoprostol. *Reproductive Health Matters* 2002;**10**:192-3.

Berer M. Medical abortion: a fact sheet. *Reprod.Health Matters.* 2005;**13**:20-4.

Chong YS, Su LL, Arulkumaran S. Misoprostol: a quarter century of use, abuse, and creative misuse. *Obstet.Gynecol.Surv.* 2004;**59**:128-40.

Dickinson JE. Late pregnancy termination within a legislated medical environment. *Aust.N.Z.J.Obstet.Gynaecol.* 2004;**44**:337-41.

Gebhardt DO. Misoprostol in a topsyturvy world. *J.Med.Ethics* 2001;**27**:205.

Weeks AD, Fiala C, Safar P. Misoprostol and the debate over off-label drug use. *BJOG.* 2005;**112**:269-72.

## Editorials/ Letters/ Opinions

Devoe L. Outcomes of second-trimester pregnancy terminations with misoprostol: Comparing 2 regimens - Discussion. *American Journal of Obstetrics and Gynecology* 2005;**193**:548-50.

Edwards. Outcomes of second-trimester pregnancy terminations with misoprostol: Comparing 2 regimens - Closing statement. *American Journal of Obstetrics and Gynecology* 2005;**193**:550.

Gebhardt DO. Misoprostol in a topsyturvy world. *J.Med.Ethics* 2001;**27**:205.

Hidar S, Jerbi M, Khairi H. Association of intravenous oxytocin to vaginal misoprostol for mid trimester labor induction. *Am.J.Obstet.Gynecol.* 2006;**195**:332-3.

Jain JK, Mishell DR. 2Nd-Trimester Abortion - Reply. *New England Journal of Medicine* 1994;**331**:1716.

Jain JK, Mishell DR, Jr. Misoprostol regimens for termination of second trimester pregnancy. *Hum.Reprod.* 2001;**16**:393.

Nuthalapaty FS, Ramsey PS. Association of intravenous oxytocin to vaginal misoprostol for mid trimester labor induction - Reply. *American Journal of Obstetrics and Gynecology* 2006;**195**:333-4.

Rosenfield A. The difficult issue of second-trimester abortion. *N.Engl.J.Med.* 1994;**331**:324-5.

Trouton K. Re: a randomized trial comparing oral misoprostol with intra-amniotic prostaglandin F2alpha for second trimester terminations. *J.Obstet.Gynaecol.Can.* 2006;**28**:189.

Wong KS. Misoprostol regimens for termination of second trimester pregnancy. *Human Reproduction* 2001;**16**:393.

Young JM. 2Nd-Trimester Abortion. *New England Journal of Medicine* 1994;**331**:1716.

Young JM. Second trimester abortion. *N.Engl.J.Med.* 1994;**331**:1716.

## Unspecified

[Anon]. Self-administration of misoprostol at 10-13 weeks of pregnancy. *Reproductive Health Matters* 2000;**8**:178-9.

Airoidi JA, O'Neill AM, Einhorn K, Hoffman M, Berghella V. Risk of uterine rupture using misoprostol in the second trimester in women with prior cesarean delivery. *Obstetrics and Gynecology* 2006;**107**:59S-60S.

Hollander D. For second-trimester abortion, women given misoprostol vaginally report the greatest satisfaction. *Perspectives on Sexual and Reproductive Health* 2004;**36**:133.

Paun MF, Bulucea C. Vaginal misoprostol is a medically sound, cost-effective option for second trimester therapeutic abortion. *Human Reproduction* 1999;**14**:255.