

# Miscarriage (2<sup>nd</sup> Trimester)

## Referenced to July 2007

### Review Articles

Griebel CP, Halvorsen J, Golemon TB, Day AA. Management of spontaneous abortion. *Am.Fam.Physician* 2005;**72**:1243-50.

Neilson JP, Hickey M, Vazquez J. Medical treatment for early fetal death (less than 24 weeks). *Cochrane.Database.Syst.Rev.* 2006;**3**:CD002253.

### Clinical Studies

#### Randomised Controlled Trials

Eng NS, Guan AC. Comparative study of intravaginal misoprostol with gemeprost as an abortifacient in second trimester missed abortion. *Aust.N.Z.J.Obstet.Gynaecol.* 1997;**37**:331-4.

Phupong V, Taneepanichskul S, Kriengsinyot R, Sriyirojana N, Blanchard K, Winikoff B. Comparative study between single dose 600 microg and repeated dose of oral misoprostol for treatment of incomplete abortion. *Contraception* 2004;**70**:307-11.

#### Observational Studies

Kutlu T, Tugrul S, Aran T, Uslu H, Eren S, Oral O. Combined administration of misoprostol in the first and second trimester missed abortion cases. *Clin.Exp.Obstet.Gynecol.* 2006;**33**:226-8.

Menakaya U, Otoide V, Omo AL, Odunsi K, Okonofua F. Experience with misoprostol in the management of missed abortion in the second trimester. *J.Obstet.Gynaecol.* 2005;**25**:583-5.

Sharma D, Singhal SR, Rani XX. Sublingual misoprostol in management of missed abortion in India. *Trop.Doct.* 2007;**37**:39-40.

**Cost-effectiveness**

////////////////////////////////////

**Pharmacology**

////////////////////////////////////

**Politics/ Licensing/ Ethics**

////////////////////////////////////

**Editorials/ Letters/ Opinions**

////////////////////////////////////

**Unspecified**

////////////////////////////////////