

Prevention of Postpartum Haemorrhage

Referenced to July 2008

Review Articles

Adams V, Miller S, Craig S, Nyima, Sonam, Droyoung *et al.* The-challenge of cross-cultural clinical trials research: Case report from the Tibetan Autonomous Region, People's Republic of China. *Medical Anthropology Quarterly* 2005;**19**:267-89.

Alfirevic Z. Blum J. Walraven G. Weeks A. Winikoff B. Prevention of postpartum hemorrhage with misoprostol. *International Journal of Gynaecology & Obstetrics*. **99** Suppl 2:S198-201, 2007 Dec.

Anderson JM, Etches D. Prevention and management of postpartum hemorrhage. *American Family Physician* 2007;**75**:875-82.

Anderson T. Prostaglandins for the prevention of PPH. *Pract.Midwife*. 2005;**8**:43-5.

Bechard dS. [Use of misoprostol in gynecology and obstetrics]. *Gynecol.Obstet.Fertil.* 2002;**30**:317-24.

Bouwmeester FW, Bolte AC, van Geijn HP. Pharmacological and surgical therapy for primary postpartum hemorrhage. *Current Pharmaceutical Design* 2005;**11**:759-73.

Brucker MC. Management of the third stage of labor: An evidence-based approach. *Journal of Midwifery & Womens Health* 2001;**46**:381-92.

Chong YS, Su LL, Arulkumaran S. Current strategies for the prevention of postpartum haemorrhage in the third stage of labour. *Curr.Opin.Obstet.Gynecol.* 2004;**16**:143-50.

de Groot ANJA, van Dongen PWJ, Vree TB, Hekster YA, van Roosmalen J. Ergot alkaloids - Current status and review of clinical pharmacology and therapeutic use compared with other oxytocics in obstetrics and gynaecology. *Drugs* 1998;**56**:523-35.

Derman RJ. The role of misoprostol for prevention of postpartum hemorrhage. *International Journal of Gynecology & Obstetrics* 2006;**94**:S131-S132.

Dildy GA. Postpartum hemorrhage: New management options. *Clinical Obstetrics and Gynecology* 2002;**45**:330-44.

El Refaey H, Rodeck C. Post-partum haemorrhage: definitions, medical and surgical management. A time for change. *Br.Med.Bull.* 2003;**67**:205-17.

Geller SE, Adams MG, Kelly PJ, Kodkany BS, Derman RJ. Postpartum hemorrhage in resource-poor settings. *Int.J.Gynaecol.Obstet.* 2006;**92**:202-11.

Goldberg AB, Greenberg MB, Darney PD. Drug therapy: Misoprostol and pregnancy. *New England Journal of Medicine* 2001;**344**:38-47.

Goldberg AB, Greenberg MB, Darney PD. Misoprostol and pregnancy. *N.Engl.J.Med.* 2001;**344**:38-47.

Gulmezoglu AM. Prostaglandins for prevention of postpartum haemorrhage. *Cochrane.Database.Syst.Rev.* 2000;CD000494.

Gulmezoglu AM, Forna F, Villar J, Hofmeyr GJ. Prostaglandins for prevention of postpartum haemorrhage. *Cochrane.Database.Syst.Rev.* 2001;CD000494.

Gulmezoglu AM, Forna F, Villar J, Hofmeyr GJ. Prostaglandins for prevention of postpartum haemorrhage. *Cochrane.Database.Syst.Rev.* 2002;CD000494.

Gulmezoglu AM, Forna F, Villar J, Hofmeyr GJ. Prostaglandins for prevention of postpartum haemorrhage. *Cochrane.Database.Syst.Rev.* 2004;CD000494.

Hemmerling A. The safety of misoprostol. *International Journal of Gynecology & Obstetrics* 2006;**94**:S149-S150.

Hofmeyr GJ. Evidence-based intrapartum care. *Best.Pract.Res.Clin.Obstet.Gynaecol.* 2005;**19**:103-15.

Jadesimi A., Okonofua FE. Tackling the unacceptable: Nigeria approves misoprostol for postpartum haemorrhage. *J.Fam.Plann.Reprod.Health Care* 2006;**32**:213-4.

Joy SD, Sanchez-Ramos L, Kaunitz AM. Misoprostol use during the third stage of labor. *Int.J.Gynaecol.Obstet.* 2003;**82**:143-52.

Lalonde A, Daviss BA, Acosta A, Herschderfer K. Postpartum hemorrhage today: ICM/FIGO initiative 2004-2006. *Int.J.Gynaecol.Obstet.* 2006;**94**:243-53.

Langenbach C. Misoprostol in preventing postpartum hemorrhage: a meta-analysis. *Int.J.Gynaecol.Obstet.* 2006;**92**:10-8.

Lazarus JV, Lalonde A. Reducing postpartum hemorrhage in Africa. *Int.J.Gynaecol.Obstet.* 2005;**88**:89-90.

Lokugamage AU, Refaey HE, Rodeck CH. Misoprostol and pregnancy: ever-increasing indications of effective usage. *Curr.Opin.Obstet.Gynecol.* 2003;**15**:513-8.

Magann EF, Lanneau GS. Third stage of labor. *Obstet.Gynecol.Clin.North Am.* 2005;**32**:323-xi.

Mategrano VA, Gabay MP. Misoprostol in the prevention of postpartum hemorrhage. *Ann.Pharmacother.* 2001;**35**:1648-52.

Mathai M, Gulmezoglu AM, Hill S. Saving women's lives: evidence-based recommendations for the prevention of postpartum haemorrhage. *Bulletin of the World Health Organization* 2007;**85**:322-3.

McCormick ML, Sanghvi HC, Kinzie B, McIntosh N. Preventing postpartum hemorrhage in low-resource settings. *Int.J.Gynaecol.Obstet.* 2002;**77**:267-75.

McDonald S. Management of the third stage of labor. *J.Midwifery Womens Health* 2007;**52**:254-61.

Miller S, Lester F, Hensleigh P. Prevention and treatment of postpartum hemorrhage: new advances for low-resource settings. *J.Midwifery Womens Health* 2004;**49**:283-92.

Moore ML. Misoprostol-is more research needed? *J.Perinat.Educ.* 2002;**11**:43-7.

Mousa HA, Walkinshaw S. Major postpartum haemorrhage. *Curr.Opin.Obstet.Gynecol.* 2001;**13**:595-603.

Peek MJ. Drugs for obstetric conditions. *Best Practice & Research in Clinical Obstetrics & Gynaecology* 2001;**15**:999-1011.

Potts M, Hemmerling A. The worldwide burden of postpartum haemorrhage: Policy development where inaction is lethal. *Int.J.Gynaecol.Obstet.* 2006;**94 Suppl 2**:S116-S121.

Prata N. A global overview of ongoing misoprostol studies. *International Journal of Gynecology & Obstetrics* 2006;**94**:S147-S148.

Ramanathan G, Arulkumaran S. Postpartum hemorrhage. *J.Obstet.Gynaecol.Can.* 2006;**28**:967-73.

Ripley DL. Uterine emergencies - Atony, inversion, and rupture. *Obstetrics and Gynecology Clinics of North America* 1999;**26**:419-+.

Rizvi F, Mackey R, Barrett T, McKenna P, Geary M. Successful reduction of massive postpartum haemorrhage by use of guidelines and staff education. *Bjog-An International Journal of Obstetrics and Gynaecology* 2004;**111**:495-8.

Sharma S, El Refaey H. Prostaglandins in the prevention and management of postpartum haemorrhage. *Best.Pract.Res.Clin.Obstet.Gynaecol.* 2003;**17**:811-23.

Song J. Use of misoprostol in obstetrics and gynecology. *Obstet.Gynecol.Surv.* 2000;**55**:503-10.

Tessier V., Pierre F. [Risk factors of postpartum hemorrhage during labor and clinical and pharmacological prevention]. *J.Gynecol.Obstet.Biol.Reprod.(Paris)* 2004;**33**:4S29-56.

Thomas J. In low-resource settings, misoprostol can reduce risk of postpartum hemorrhage. *International Family Planning Perspectives* 2007;**33**:39-40.

Tsu VD, Langer A, Aldrich T. Postpartum hemorrhage in developing countries: is the public health community using the right tools? *Int.J.Gynaecol.Obstet.* 2004;**85 Suppl 1**:S42-S51.

Villar J, Gulmezoglu AM, Hofmeyr GJ, Forna F. Systematic review of randomized controlled trials of misoprostol to prevent postpartum hemorrhage. *Obstet.Gynecol.* 2002;**100**:1301-12.

Walder J. Misoprostol: preventing postpartum haemorrhage. *Mod.Midwife.* 1997;**7**:23-7.

Zust S, Hosli I, Surbek D, Holzgreve W. Side effects of Misoprostol therapy during pregnancy. *Zeitschrift fur Geburtshilfe und Neonatologie* 2001;**205**:43-8.

Zust S, Hosli I, Surbek D, Holzgreve W. [Adverse effects of misoprostol in pregnancy]. *Z.Geburtshilfe Neonatol.* 2001;**205**:43-8.

Clinical Studies

Randomised Controlled Trials

Acharya G, Al Sammarai MT, Patel N, Al Habib A, Kiserud T. A randomized, controlled trial comparing effect of oral misoprostol and intravenous syntocinon on intra-operative blood loss during cesarean section. *Acta Obstet.Gynecol.Scand.* 2001;**80**:245-50.

Amant F, Spitz B, Timmerman D, Corremans A, Van Assche FA. Misoprostol compared with methylergometrine for the prevention of postpartum haemorrhage: a double-blind randomised trial. *Br.J.Obstet.Gynaecol.* 1999;**106**:1066-70.

Bamigboye AA, Hofmeyr GJ, Merrell DA. Rectal misoprostol in the prevention of postpartum hemorrhage: a placebo-controlled trial. *Am.J.Obstet.Gynecol.* 1998;**179**:1043-6.

Bamigboye AA, Merrell DA, Hofmeyr GJ, Mitchell R. Randomized comparison of rectal misoprostol with Syntometrine for management of third stage of labor. *Acta Obstet.Gynecol.Scand.* 1998;**77**:178-81.

- Baskett TF, Persad VL, Clough HJ, Young DC. Misoprostol versus oxytocin for the reduction of postpartum blood loss. *Int.J.Gynaecol.Obstet.* 2007;**97**:2-5.
- Benchimol M, Gondry J, Mention JE, Gagneur O, Boulanger JC. [Role of misoprostol in the delivery outcome]. *J.Gynecol.Obstet.Biol.Reprod.(Paris)* 2001;**30**:576-83.
- Bhullar A, Carlan SJ, Hamm J, Lamberty N, White L, Richichi K. Buccal misoprostol to decrease blood loss after vaginal delivery: a randomized trial. *Obstet.Gynecol.* 2004;**104**:1282-8.
- Bugalho A, Daniel A, Faundes A, Cunha M. Misoprostol for prevention of postpartum hemorrhage. *Int.J.Gynaecol.Obstet.* 2001;**73**:1-6.
- Caliskan E, Meydanli MM, Dilbaz B, Aykan B, Sonmezer M, Haberal A. Is rectal misoprostol really effective in the treatment of third stage of labor? A randomized controlled trial. *Am.J.Obstet.Gynecol.* 2002;**187**:1038-45.
- Caliskan E, Dilbaz B, Meydanli MM, Ozturk N, Narin MA, Haberal A. Oral misoprostol for the third stage of labor: a randomized controlled trial. *Obstet.Gynecol.* 2003;**101**:921-8.
- Chakraborty H, Bartz J, Goudar SS. Estimating the effect of unplanned interventions in the randomized misoprostol trial: A mixed model analysis. *International Journal of Gynecology & Obstetrics* 2006;**94**:S136.
- Chandhiok N, Dhillon BS, Datey S, Mathur A, Saxena NC. Oral misoprostol for prevention of postpartum hemorrhage by paramedical workers in India. *Int.J.Gynaecol.Obstet.* 2006;**92**:170-5.
- Chong YS, Chua S, Shen L, Arulkumaran S. Does the route of administration of misoprostol make a difference? The uterotonic effect and side effects of misoprostol given by different routes after vaginal delivery. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2004;**113**:191-8.
- Cook CM, Spurrett B, Murray H. A randomized clinical trial comparing oral misoprostol with synthetic oxytocin or syntometrine in the third stage of labour. *Aust.N.Z.J.Obstet.Gynaecol.* 1999;**39**:414-9.
- Derman RJ, Kodkany BS, Goudar SS, Geller SE, Naik VA, Bellad MB *et al.* Oral misoprostol in preventing postpartum haemorrhage in resource-poor communities: a randomised controlled trial. *Lancet* 2006;**368**:1248-53.
- El Refaey H, Nooh R, O'Brien P, Abdalla M, Geary M, Walder J *et al.* The misoprostol third stage of labour study: a randomised controlled comparison between orally administered misoprostol and standard management. *BJOG.* 2000;**107**:1104-10.

Enakpene CA, Morhason-Bello IO, Enakpene EO, Arowojolu AO, Omigbodun AO. Oral misoprostol for the prevention of primary post-partum hemorrhage during third stage of labor. *Journal of Obstetrics & Gynaecology Research*. **33**(6):810-7, 2007 Dec.

Garg P, Batra S, Gandhi G. Oral misoprostol versus injectable methylergometrine in management of the third stage of labor. *Int.J.Gynaecol.Obstet*. 2005;**91**:160-1.

Gerstenfeld TS, Wing DA. Rectal misoprostol versus intravenous oxytocin for the prevention of postpartum hemorrhage after vaginal delivery. *Am.J.Obstet.Gynecol*. 2001;**185**:878-82.

Gulmezoglu AM, Villar J, Ngoc NT, Piaggio G, Carroli G, Adetoro L *et al*. WHO multicentre randomised trial of misoprostol in the management of the third stage of labour. *Lancet* 2001;**358**:689-95.

Gupta B, Jain V, Aggarwal N. Rectal misoprostol versus oxytocin in the prevention of postpartum hemorrhage - A pilot study. *International Journal of Gynecology & Obstetrics* 2006;**94**:S139-S140.

Hamm J, Russell Z, Botha T, Carlan SJ, Richichi K. Buccal misoprostol to prevent hemorrhage at cesarean delivery: a randomized study. *Am.J.Obstet.Gynecol*. 2005;**192**:1404-6.

Hofmeyr GJ, Nikodem VC, de Jager M, Gelbart BR. A randomised placebo controlled trial of oral misoprostol in the third stage of labour. *Br.J.Obstet.Gynaecol*. 1998;**105**:971-5.

Hofmeyr GJ, Nikodem VC, de Jager M, Drakely A. Side-effects of oral misoprostol in the third stage of labour--a randomised placebo-controlled trial. *S.Afr.Med.J*. 2001;**91**:432-5.

Hoj L, Cardoso P, Nielsen BB, Hvidman L, Nielsen J, Aaby P. Effect of sublingual misoprostol on severe postpartum haemorrhage in a primary health centre in Guinea-Bissau: randomised double blind clinical trial. *BMJ* 2005;**331**:723.

Karkanis SG, Caloia D, Salenieks ME, Kingdom J, Walker M, Meffe F *et al*. Randomized controlled trial of rectal misoprostol versus oxytocin in third stage management. *J.Obstet.Gynaecol.Can*. 2002;**24**:149-54.

Khan RU, El Refaey H. Pharmacokinetics and adverse-effect profile of rectally administered misoprostol in the third stage of labor. *Obstet.Gynecol*. 2003;**101**:968-74.

Kodkany BS, Derman RJ, Goudar SS, Geller SE, Edlavitch SA, Naik VA *et al*. Initiating a novel therapy in preventing postpartum hemorrhage in rural India: a joint collaboration between the United States and India. *Int.J.Fertil.Womens Med*. 2004;**49**:91-6.

Kodkany BS, Goudar SS, Derman RJ. The efficacy of oral misoprostol in preventing postpartum hemorrhage in a community setting: A randomized double-blind placebo-controlled trial. *International Journal of Gynecology & Obstetrics* 2006;**94**:S141-S142.

Kundodyiwa TW, Majoko F, Rusakaniko S. Misoprostol versus oxytocin in the third stage of labor. *Int.J.Gynaecol.Obstet.* 2001;**75**:235-41.

Lam H, Tang OS, Lee CP, Ho PC. A pilot-randomized comparison of sublingual misoprostol with syntometrine on the blood loss in third stage of labor. *Acta Obstet.Gynecol.Scand.* 2004;**83**:647-50.

Lapaire O, Schneider MC, Stotz M, Surbek DV, Holzgreve W, Hoesli IM. Oral misoprostol vs. intravenous oxytocin in reducing blood loss after emergency cesarean delivery. *Int.J.Gynaecol.Obstet.* 2006;**95**:2-7.

Lokugamage AU, Paine M, Bassaw-Balroop K, Sullivan KR, Refaey HE, Rodeck CH. Active management of the third stage at caesarean section: a randomised controlled trial of misoprostol versus syntocinon. *Aust.N.Z.J.Obstet.Gynaecol.* 2001;**41**:411-4.

Lumbiganon P, Hofmeyr J, Gulmezoglu AM, Pinol A, Villar J. Misoprostol dose-related shivering and pyrexia in the third stage of labour. WHO Collaborative Trial of Misoprostol in the Management of the Third Stage of Labour. *Br.J.Obstet.Gynaecol.* 1999;**106**:304-8.

Mahajan NN, Mahajan K, Soni R. A Double-Blind Randomized Controlled Trial of Oral Misoprostol and Intramuscular Syntometrine in the Management of the Third Stage of Labor. *Gynecol.Obstet.Invest* 2007;**64**:82.

Nellore V, Mittal S, Dadhwal V. Rectal misoprostol vs. 15-methyl prostaglandin F₂alpha for the prevention of postpartum hemorrhage. *Int.J.Gynaecol.Obstet.* 2006;**94**:45-6.

Ng PS, Chan AS, Sin WK, Tang LC, Cheung KB, Yuen PM. A multicentre randomized controlled trial of oral misoprostol and i.m. syntometrine in the management of the third stage of labour. *Hum.Reprod.* 2001;**16**:31-5.

Ng PS, Lai CY, Sahota DS, Yuen PM. A double-blind randomized controlled trial of oral misoprostol and intramuscular syntometrine in the management of the third stage of labor. *Gynecol.Obstet.Invest* 2007;**63**:55-60.

Nielsen BB, Hoj L, Hvidman LE, Nielsen J, Cardoso P, Aaby P. [Reduced postpartum bleeding after treatment with sublingual misoprostol: a randomized double-blind clinical study in a developing country--secondary publication)]. *Ugeskr.Laeger* 2006;**168**:1341-3.

Oboro VO, Tabowe TO. A randomised controlled trial of misoprostol versus oxytocin in the active management of the third stage of labour. *J.Obstet.Gynaecol.* 2003;**23**:13-6.

- Ozkaya O, Sezik M, Kaya H, Desdicioglu R, Dittrich R. Placebo-controlled randomized comparison of vaginal with rectal misoprostol in the prevention of postpartum hemorrhage. *J.Obstet.Gynaecol.Res.* 2005;**31**:389-93.
- Parsons SM, Walley RL, Crane JM, Matthews K, Hutchens D. Oral misoprostol versus oxytocin in the management of the third stage of labour. *J.Obstet.Gynaecol.Can.* 2006;**28**:20-6.
- Piaggio G, Elbourne D, Schulz KF, Villar J, Pinol AP, Gulmezoglu AM. The reporting of methods for reducing and detecting bias: an example from the WHO Misoprostol Third Stage of Labour equivalence randomised controlled trial. *BMC.Med.Res.Methodol.* 2003;**3**:19.
- Saleem S. Efficacy of dinoprostone, intracervical foleys and misoprostol in labor induction. *J.Coll.Physicians Surg.Pak.* 2006;**16**:276-9.
- Schaff EA, DiCenzo R, Fielding SL. Comparison of misoprostol plasma concentrations following buccal and sublingual administration. *Contraception* 2005;**71**:22-5.
- Surbek DV, Fehr PM, Hosli I, Holzgreve W. Oral misoprostol for third stage of labor: a randomized placebo-controlled trial. *Obstet.Gynecol.* 1999;**94**:255-8.
- Tudor C, Miller S, Nyima, Sonam, Droyoung, Varner M. Preliminary progress report: Randomized double-blind trial of Zhi Byed 11, a Tibetan traditional medicine, versus misoprostol to prevent postpartum hemorrhage in Lhasa, Tibet. *International Journal of Gynecology & Obstetrics* 2006;**94**:S145-S146.
- Verma P, Aggarwal N, Jain V, Suri V. A double-blind randomized controlled trial to compare sublingual misoprostol with methylergometrine for prevention of postpartum hemorrhage. *International Journal of Gynecology & Obstetrics* 2006;**94**:S137-S138.
- Vimala N, Mittal S, Kumar S, Dadhwal V, Mehta S. Sublingual misoprostol versus methylergometrine for active management of the third stage of labor. *Int.J.Gynaecol.Obstet.* 2004;**87**:1-5.
- Vimala N, Mittal S, Kumar S. Sublingual misoprostol versus oxytocin infusion to reduce blood loss at cesarean section. *Int.J.Gynaecol.Obstet.* 2006;**92**:106-10.
- Vogel D, Burkhardt T, Rentsch K, Schweer H, Watzer B, Zimmermann R *et al.* Misoprostol versus methylergometrine: pharmacokinetics in human milk. *Am.J.Obstet.Gynecol.* 2004;**191**:2168-73.
- Walley RL, Wilson JB, Crane JM, Matthews K, Sawyer E, Hutchens D. A double-blind placebo controlled randomised trial of misoprostol and oxytocin in the management of the third stage of labour. *BJOG.* 2000;**107**:1111-5.

Wright JM, Newton W. Is rectal misoprostol as effective as oxytocin in preventing postpartum hemorrhage? *J.Fam.Pract.* 2003;**52**:281-2.

Zachariah ES, Naidu M, Seshadri L. Oral misoprostol in the third stage of labor. *Int.J.Gynaecol.Obstet.* 2006;**92**:23-6.

Zhao Y, Li X, Peng Y. [Clinical study on reduction of postpartum bleeding in cesarean section by misoprostol]. *Zhonghua Fu Chan Ke.Za Zhi.* 1998;**33**:403-5.

Observational Studies

Abdel-Aleem H, Villar J, Gulmezoglu AM, Mostafa SA, Youssef AA, Shokry M *et al.* The pharmacokinetics of the prostaglandin E1 analogue misoprostol in plasma and colostrum after postpartum oral administration. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2003;**108**:25-8.

Andolina KL, Tolosa JE, Monzo JM, Roberts NS, Daly S. Oral misoprostol is rapidly absorbed in postpartum women at term. *J.Matern.Fetal Neonatal Med.* 2003;**14**:229-32.

Bjornerem A, Acharya G, Oian P, Maltau JM. [Postpartum hemorrhage--prophylaxis and treatment in Norway]. *Tidsskr.Nor Laegeforen.* 2002;**122**:2536-7.

Chandra S, Persad V, Young D, Baskett T. A preliminary study of cutaneous blood flow associated with postpartum use of oral misoprostol. *J.Obstet.Gynaecol.Can.* 2004;**26**:1073-6.

Chong YS, Chua S, El Refaey H, Choo WL, Chanrachakul B, Tai BC *et al.* Postpartum intrauterine pressure studies of the uterotonic effect of oral misoprostol and intramuscular syntometrine. *BJOG.* 2001;**108**:41-7.

Choo WL, Chua S, Chong YS, Vanaja K, Oei PL, Ho LM *et al.* Correlation of change in uterine activity to blood loss in the third stage of labour. *Gynecol.Obstet.Invest* 1998;**46**:178-80.

Clark S, Blum J, Blanchard K, Galvao L, Fletcher H, Winikoff B. Misoprostol use in obstetrics and gynecology in Brazil, Jamaica, and the United States. *Int.J.Gynaecol.Obstet.* 2002;**76**:65-74.

de Vaate AB, Coleman R, Manneh H, Walraven G. Knowledge, attitudes and practices of trained traditional birth attendants in the Gambia in the prevention, recognition and management of postpartum haemorrhage. *Midwifery* 2002;**18**:3-11.

Diab KM, Ramy AR, Yehia MA. The use of rectal misoprostol as active pharmacological management of the third stage of labor. *J.Obstet.Gynaecol.Res.* 1999;**25**:327-32.

El Refaey H, O'Brien P, Morafa W, Walder J, Rodeck C. Use of oral misoprostol in the prevention of postpartum haemorrhage. *Br.J.Obstet.Gynaecol.* 1997;**104**:336-9.

Prata N, Hamza S, Gypson R, Nada K, Vahidnia F, Potts M. Misoprostol and active management of the third stage of labor. *Int.J.Gynaecol.Obstet.* 2006;**94**:149-55.

Quiroga DR, Esparza AM, Batiza R, V, Coronado LO, Hernandez AS, Martinez CJ. [Vaginal misoprostol in the prevention of postpartum hemorrhage]. *Ginecol.Obstet.Mex.* 2002;**70**:572-5.

Rajbhandari S, Pun A, Hodgins S, Rajendra PK. Prevention of postpartum haemorrhage at homebirth with use of misoprostol in Banke District, Nepal. *International Journal of Gynecology & Obstetrics* 2006;**94**:S143-S144.

Sina ED. The reduction in maternal mortality in Chile, 1990-200. *Revista Panamericana de Salud Publica-Pan American Journal of Public Health* 2004;**15**:326-30.

Cost-effectiveness

////////////////////////////////////

Pharmacology

Abdel-Aleem H, Villar J, Gulmezoglu AM, Mostafa SA, Youssef AA, Shokry M *et al.* The pharmacokinetics of the prostaglandin E1 analogue misoprostol in plasma and colostrum after postpartum oral administration. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2003;**108**:25-8.

Andolina KL, Tolosa JE, Monzo JM, Roberts NS, Daly S. Oral misoprostol is rapidly absorbed in postpartum women at term. *J.Matern.Fetal Neonatal Med.* 2003;**14**:229-32.

Chandra S, Persad V, Young D, Baskett T. A preliminary study of cutaneous blood flow associated with postpartum use of oral misoprostol. *J.Obstet.Gynaecol.Can.* 2004;**26**:1073-6.

Chong YS, Chua S, Arulkumaran S. Severe hyperthermia following oral misoprostol in the immediate postpartum period. *Obstet.Gynecol.* 1997;**90**:703-4.

Goldberg AB, Greenberg MB, Darney PD. Misoprostol and pregnancy. *N.Engl.J.Med.* 2001;**344**:38-47.

Htay TT. Making pregnancy safer in Myanmar: introducing misoprostol to prevent post-partum haemorrhage as part of active management of the third stage of labour. *Reproductive Health Matters.* **15**(30):214-5, 2007 Nov.

Hemmerling A. The safety of misoprostol. *International Journal of Gynecology & Obstetrics* 2006;**94**:S149-S150.

Khan RU, El Refaey H. Pharmacokinetics and adverse-effect profile of rectally administered misoprostol in the third stage of labor. *Obstet.Gynecol.* 2003;**101**:968-74.

Lumbiganon P, Villar J, Piaggio G, Gulmezoglu AM, Adetoro L, Carroli G. Side effects of oral misoprostol during the first 24 hours after administration in the third stage of labour. *BJOG.* 2002;**109**:1222-6.

Schaff EA, DiCenzo R, Fielding SL. Comparison of misoprostol plasma concentrations following buccal and sublingual administration. *Contraception* 2005;**71**:22-5.

Vogel D, Burkhardt T, Rentsch K, Schweer H, Watzer B, Zimmermann R *et al.* Misoprostol versus methylergometrine: pharmacokinetics in human milk. *Am.J.Obstet.Gynecol.* 2004;**191**:2168-73.

Zust S, Hosli I, Surbek D, Holzgreve W. Side effects of Misoprostol therapy during pregnancy. *Zeitschrift fur Geburtshilfe und Neonatologie* 2001;**205**:43-8.

Zust S, Hosli I, Surbek D, Holzgreve W. [Adverse effects of misoprostol in pregnancy]. *Z.Geburtshilfe Neonatol.* 2001;**205**:43-8.

Politics/ Licensing/ Ethics

Campbell M, Holden M. Global availability of misoprostol. *International Journal of Gynecology & Obstetrics* 2006;**94**:S151-S152.

Chong YS, Su LL, Arulkumaran S. Misoprostol: a quarter century of use, abuse, and creative misuse. *Obstet.Gynecol.Surv.* 2004;**59**:128-40.

Darney PD. Misoprostol: a boon to safe motherhood...or not? *Lancet* 2001;**358**:682-3.

Gebhardt DO. Misoprostol in a topsyturvy world. *J.Med.Ethics* 2001;**27**:205.

Potts M, Campbell M. Three meetings and fewer funerals--misoprostol in postpartum haemorrhage. *Lancet* 2004;**364**:1110-1.

Weeks AD, Fiala C, Safar P. Misoprostol and the debate over off-label drug use. *BJOG*. 2005;**112**:269-72.

Editorials/ Letters/ Opinions

Amant F. The misoprostal third stage study: a randomised controlled comparison between orally administered misoprostol and standard management. A double-blind placebo controlled randomised trial of misoprostol and oxytocin in the management of the third stage labour. *BJOG*. 2001;**108**:338.

Carpenter JP. Misoprostol for prevention of postpartum haemorrhage in developing countries. *Trop.Doct.* 2003;**33**:122.

Chong YS, Su LL. Misoprostol for preventing PPH: some lessons learned. *Lancet* 2006;**368**:1216-8.

El Refaey H, O'Brien P, Morafa W, Walder J, Rodeck C. Misoprostol for third stage of labour. *Lancet* 1996;**347**:1257.

El Refaey H. Use of misoprostol in third stage of labour. *Lancet* 2002;**359**:707-8.

Gebhardt DO. Misoprostol in a topsyturvy world. *J.Med.Ethics* 2001;**27**:205.

Hofmeyr GJ. Misoprostol in obstetrics and gynaecology--unregistered, dangerous and essential. *S.Afr.Med.J.* 1998;**88**:535-6.

Hofmeyr GJ, Bamigboye AA. Rectal misoprostol in the prevention of postpartum hemorrhage - Reply. *American Journal of Obstetrics and Gynecology* 1999;**180**:1601-2.

Khan R, Sharma S. Use of misoprostol in third stage of labour. *Lancet* 2002;**359**:708-9.

Lee N. Postpartum hemorrhage. *Am.J.Nurs.* 2007;**107**:15.

Mahajan NN, Mahajan K, Soni R. A double-blind randomized controlled trial of oral misoprostol and intramuscular syntometrine in the management of the third stage of labor. *Gynecologic & Obstetric Investigation*. **64**(2):82; author reply 83, 2007.

O'Brien P, Lokugamage AU, Guillebaud J, Rodeck CH. Use of misoprostol in third stage of labour. *Lancet* 2002;**359**:708-10.

Olufowobi O, Afnan M, Sorinola O, Papaioannou S, Sharif K. Re: A randomized study comparing rectally administered misoprostol versus syntometrine combined with an oxytocin infusion for the cessation of primary postpartum hemorrhage. *Acta Obstetricia et Gynecologica Scandinavica* 2002;**81**:993.

Ramsey PS, Ramin KD, Bradley SS. Rectal misoprostol in the prevention of postpartum hemorrhage. *Am.J.Obstet.Gynecol.* 1999;**180**:1601-2.

Sciarra JJ. Announcing the winner of the IJGO Prize Paper Award for 2001. *Int.J.Gynaecol.Obstet.* 2002;**77**:105-7.

Shannon C., Winikoff B. Use of misoprostol in third stage of labour. *Lancet* 2002;**359**:709-10.

Villar J, Guimezoglu AM, Hofmeyr GJ, Cheng LN, Elbourne D. Use of misoprostol in third stage of labour - Reply. *Lancet* 2002;**359**:709-10.

Weeks A. Oral misoprostol for postpartum haemorrhage. *Lancet* 2006;**368**:2123.

Unspecified

Baskett TF, Persad V, Clough H, Young D. Prophylactic use of misoprostol in the third stage of labor. *Obstetrics and Gynecology* 2005;**105**:39S.