

Induction of Labour

Referenced to July 2008

Review Articles

Committee opinion. Induction of labor for vaginal birth after cesarean delivery. *Obstet.Gynecol.* 2002;**99**:679-80.

SOGC clinical practice guidelines. Guidelines for vaginal birth after previous caesarean birth. Number 155 (Replaces guideline Number 147), February 2005. *Int.J.Gynaecol.Obstet.* 2005;**89**:319-31.

ACOG Committee Opinion No. 342: induction of labor for vaginal birth after cesarean delivery. *Obstet.Gynecol.* 2006;**108**:465-8.

[Anon]. Urgent need for standardised misoprostol regimens. *Reproductive Health Matters* 2002;**10**:202.

[Anon]. Vaginal birth after previous cesarean section. *Journal of Midwifery & Womens Health* 2004;**49**:68-75.

Alfirevic Z, Howarth G, Gaussmann A. Oral misoprostol for induction of labour with a viable fetus. *Cochrane.Database.Syst.Rev.* 2000;CD001338.

Alfirevic Z. Oral misoprostol for induction of labour. *Cochrane.Database.Syst.Rev.* 2000;CD001338.

Alfirevic Z. Oral misoprostol for induction of labour. *Cochrane.Database.Syst.Rev.* 2001;CD001338.

Alfirevic Z, Weeks A. Oral misoprostol for induction of labour. *Cochrane.Database.Syst.Rev.* 2006;CD001338.

Apgar BS. Current trends in cervical ripening and labor induction. *Am.Fam.Physician* 1999;**60**:418-20.

Bartusevicius A, Barcaite E, Nadisauskiene R. Oral, vaginal and sublingual misoprostol for induction of labor. *Int.J.Gynaecol.Obstet.* 2005;**91**:2-9.

Bauer TA, Brown DL, Chai LK. Vaginal misoprostol for term labor induction. *Ann.Pharmacother.* 1997;**31**:1391-3.

Baulieu EE. The combined use of prostaglandin and antiprogestin in human fertility control. *Adv.Prostaglandin Thromboxane Leukot.Res.* 1995;**23**:55-62.

- Baxley EG. Labor induction: A decade of change. *American Family Physician* 2003;**67**:2076-+.
- Bechard dS. [Use of misoprostol in gynecology and obstetrics]. *Gynecol.Obstet.Fertil.* 2002;**30**:317-24.
- Blanchard K, Clark S, Winikoff B, Gaines G, Kabani G, Shannon C. Misoprostol for women's health: a review. *Obstet.Gynecol.* 2002;**99**:316-32.
- Boulvain M, Kelly A, Lohse C, Stan C, Irion O. Mechanical methods for induction of labour. *Cochrane.Database.Syst.Rev.* 2001;CD001233.
- Brennand J, Greer I. Induction of labour: new horizons. *Hosp.Med.* 1998;**59**:856-60.
- Bugnon A, Paniagua AE, Postiglione G, Lardizabal JL. [Induction of labor with misoprostol]. *Ginecol.Obstet.Mex.* 1994;**62**:407-14.
- Capilla MC, Bermejo VT. [Efficacy and safety of misoprostol in obstetrics]. *Farm.Hosp.* 2005;**29**:177-84.
- Carr CA, Burkhardt P, Avery M. Vaginal birth after cesarean birth: A national survey of US midwifery practice. *Journal of Midwifery & Womens Health* 2002;**47**:347-52.
- Crane J, Butler B, Young D, Hannah M. Misoprostol for labor induction in women at term with intact membranes and an unfavorable cervix: A systematic review. *American Journal of Obstetrics and Gynecology* 2005;**193**:S40.
- Crane JM, Bennett KA. A META-ANALYSIS OF CONTROLLED-RELEASE PROSTAGLANDIN FOR CERVICAL RIPENING AND LABOUR INDUCTION. *J.SOGC.* 2000;**22**:692-8.
- Crane JM, Young DC. Induction of labour with a favourable cervix and/or pre-labour rupture of membranes. *Best.Pract.Res.Clin.Obstet.Gynaecol.* 2003;**17**:795-809.
- Crane JM, Butler B, Young DC, Hannah ME. Misoprostol compared with prostaglandin E2 for labour induction in women at term with intact membranes and unfavourable cervix: a systematic review. *BJOG.* 2006;**113**:1366-76.
- Dare MR, Middleton P, Crowther CA, Flenady VJ, Varatharaju B. Planned early birth versus expectant management (waiting) for prelabour rupture of membranes at term (37 weeks or more). *Cochrane Database of Systematic Reviews* 2006.
- Dodd J, Crowther C. Induction of labour for women with a previous Caesarean birth: a systematic review of the literature. *Aust.N.Z.J.Obstet.Gynaecol.* 2004;**44**:392-5.
- Duff P. Premature rupture of the membranes in term patients. *Seminars in Perinatology* 1996;**20**:401-8.

El Refaey H, Jauniaux E. Methods of induction of labour. *Curr. Opin. Obstet. Gynecol.* 1997;**9**:375-8.

Enoch J. Misoprostol (Cytotec): a new method of inducing labor. *Midwifery Today Int. Midwife.* 1999;22-30.

Ginath S, Zakut HV. Misoprostol--for cervical ripening? *Eur. J. Obstet. Gynecol. Reprod. Biol.* 2001;**99**:152-3.

Ginath S, Zakut HV. Misoprostol - for cervical ripening? *European Journal of Obstetrics Gynecology and Reproductive Biology* 2001;**99**:152-3.

Goldberg AB, Greenberg MB, Darney PD. Drug therapy: Misoprostol and pregnancy. *New England Journal of Medicine* 2001;**344**:38-47.

Goldberg AB, Greenberg MB, Darney PD. Misoprostol and pregnancy. *N. Engl. J. Med.* 2001;**344**:38-47.

Goldberg AB, Wing DA. Induction of labor: the misoprostol controversy. *J. Midwifery Womens Health* 2003;**48**:244-8.

Hadi H. Cervical ripening and labor induction: Clinical guidelines. *Clinical Obstetrics and Gynecology* 2000;**43**:524-36.

Harman JH, Jr., Kim A. Current trends in cervical ripening and labor induction. *Am. Fam. Physician* 1999;**60**:477-84.

Hemmerling A. The safety of misoprostol. *International Journal of Gynecology & Obstetrics* 2006;**94**:S149-S150.

Hofmeyr GJ, Gulmezoglu AM. Vaginal misoprostol for cervical ripening and labour induction in late pregnancy. *Birth* 1999;**26**:264.

Hofmeyr GJ, Gulmezoglu AM, Alfirevic Z. Misoprostol for induction of labour: a systematic review. *Br. J. Obstet. Gynaecol.* 1999;**106**:798-803.

Hofmeyr GJ, Gulmezoglu AM. Vaginal misoprostol for cervical ripening and labour induction in late pregnancy. *Cochrane. Database. Syst. Rev.* 2000;CD000941.

Hofmeyr GJ. Induction of labour with misoprostol. *Curr. Opin. Obstet. Gynecol.* 2001;**13**:577-81.

Hofmeyr GJ, Gulmezoglu AM. Vaginal misoprostol for cervical ripening and induction of labour. *Cochrane. Database. Syst. Rev.* 2001;CD000941.

Hofmeyr GJ, Gulmezoglu AM. Vaginal misoprostol for cervical ripening and induction of labour. *Cochrane. Database. Syst. Rev.* 2003;CD000941.

Hughes EG, Kelly AJ, Kavanagh J. Dinoprostone vaginal insert for cervical ripening and labor induction: a meta-analysis. *Obstet.Gynecol.* 2001;**97**:847-55.

Justus HG. Induction of labour with an unfavourable cervix. *Best.Pract.Res.Clin.Obstet.Gynaecol.* 2003;**17**:777-94.

Kelly AJ, Kavanagh J, Thomas J. Vaginal prostaglandin (PGE2 and PGF2a) for induction of labour at term. *Cochrane.Database.Syst.Rev.* 2003;CD003101.

King TL. Clinical controversies: applying evidence in practice. *J.Midwifery Womens Health* 2003;**48**:243.

Li XM, Wan J, Xu CF, Zhang Y, Fang L, Shi ZJ *et al.* Misoprostol in labour induction of term pregnancy: a meta-analysis. *Chin Med.J.(Engl.)* 2004;**117**:449-52.

Lin MG, Nuthalapaty F, Carver A, Case A, Ramsey P. Misoprostol compared with placebo for labor induction in women with PROM: A meta-analysis. *Obstetrics and Gynecology* 2005;**105**:79S.

Lin MG, Nuthalapaty FS, Carver AR, Case AS, Ramsey PS. Misoprostol for labor induction in women with term premature rupture of membranes: a meta-analysis. *Obstet.Gynecol.* 2005;**106**:593-601.

Lokugamage AU, Refaey HE, Rodeck CH. Misoprostol and pregnancy: ever-increasing indications of effective usage. *Curr.Opin.Obstet.Gynecol.* 2003;**15**:513-8.

Mauldin JG, Newman RB. Prior cesarean: a contraindication to labor induction? *Clin.Obstet.Gynecol.* 2006;**49**:684-97.

McDonagh MS, Osterweil P, Guise JM. The benefits and risks of inducing labour in patients with prior caesarean delivery: a systematic review. *Bjog-An International Journal of Obstetrics and Gynaecology* 2005;**112**:1007-15.

McGill J, Shetty A. Mifepristone and misoprostol in the induction of labor at term. *International Journal of Gynaecology & Obstetrics.* **96**(2):80-4, 2007 Feb

McLean MT. Labor induction with misoprostol: a midwife's perspective. *Midwifery Today Int.Midwife.* 1999;28.

Moore ML. Misoprostol-is more research needed? *J.Perinat.Educ.* 2002;**11**:43-7.

Morey SS. ACOG develops guidelines for induction of labor. *Am.Fam.Physician* 2000;**62**:445.

Mozurkewich E. Management of premature rupture of membranes at term: An evidence-based approach. *Clinical Obstetrics and Gynecology* 1999;**42**:749-56.

- Mozurkewich E. Prelabor rupture of membranes at term: induction techniques. *Clin.Obstet.Gynecol.* 2006;**49**:672-83.
- Muzonzini G., Hofmeyr GJ. Buccal or sublingual misoprostol for cervical ripening and induction of labour. *Cochrane.Database.Syst.Rev.* 2004;CD004221.
- O'Brien-Abel N. Uterine rupture during VBAC trial of labor: Risk factors and fetal response. *Journal of Midwifery & Womens Health* 2003;**48**:249-57.
- Peek MJ. Drugs for obstetric conditions. *Best Practice & Research in Clinical Obstetrics & Gynaecology* 2001;**15**:999-1011.
- Poma PA. Cervical ripening - A review and recommendations for clinical practice. *Journal of Reproductive Medicine* 1999;**44**:657-68.
- Poma PA. Cervical ripening. A review and recommendations for clinical practice. *J.Reprod.Med.* 1999;**44**:657-68.
- Ramsey PS, Ramin KD, Ramin SM. Labor induction. *Current Opinion in Obstetrics & Gynecology* 2000;**12**:463-73.
- Rayburn WF. Preinduction cervical ripening: basis and methods of current practice. *Obstet.Gynecol.Surv.* 2002;**57**:683-92.
- Reichler A, Romem Y, Divon MY. Induction of labor. *Curr.Opin.Obstet.Gynecol.* 1995;**7**:432-6.
- Roberts LM, Homer CS, Davis GK, Miller TD. Misoprostol to induce labour: a review of its use in a NSW hospital. *Australian & New Zealand Journal of Obstetrics & Gynaecology.* **47**(4):291-6, 2007 Aug.
- Rosen R. Planned early birth vs. expectant management for PROM. *Am.Fam.Physician* 2006;**74**:79-80.
- Sanchez-Ramos L, Kaunitz AM, Wears RL, Delke I, Gaudier FL. Misoprostol for cervical ripening and labor induction: a meta-analysis. *Obstet.Gynecol.* 1997;**89**:633-42.
- Sanchez-Ramos L, Kaunitz AM, Delke I, Gaudier FL. Cervical ripening and labor induction with a controlled-release dinoprostone vaginal insert: A meta-analysis. *Obstetrics and Gynecology* 1999;**94**:878-83.
- Sanchez-Ramos L, Kaunitz AM. Misoprostol for cervical ripening and labor induction: a systematic review of the literature. *Clin.Obstet.Gynecol.* 2000;**43**:475-88.
- Sanchez-Ramos L, Kaunitz AM, Delke I. Labor induction with 25 microg versus 50 microg intravaginal misoprostol: a systematic review. *Obstet.Gynecol.* 2002;**99**:145-51.

Sanchez-Ramos L, Hsieh E. Pharmacologic methods for cervical ripening and labor induction. *Curr.Womens Health Rep.* 2003;**3**:55-60.

Sitruk-Ware R. Mifepristone and misoprostol sequential regimen side effects, complications and safety. *Contraception* 2006;**74**:48-55.

Song J. Use of misoprostol in obstetrics and gynecology. *Obstet.Gynecol.Surv.* 2000;**55**:503-10.

Summers L. Methods of cervical ripening and labor induction. *J.Nurse Midwifery* 1997;**42**:71-85.

Surbek DV, Hosli I, Holzgreve W. [Current aspects of labor induction]. *Ther.Umsch.* 2002;**59**:650-9.

Taylor SJ, Armour CL. Cost implications in the management of induction of labour. *Pharmacoeconomics* 1997;**12**:547-54.

Templeton A. Misoprostol for all? *Br.J.Obstet.Gynaecol.* 1998;**105**:937-9.

Tenore JL. Methods for cervical ripening and induction of labor. *Am.Fam.Physician* 2003;**67**:2123-8.

Tinelli A, Tinelli R, Tinelli FG. [Induction of labour: which method to use?]. *Minerva Ginecol.* 2003;**55**:463-82.

Votipka JR. Misoprostol for cervical ripening and labor induction. *J.Fam.Pract.* 1997;**45**:20.

Weaver SP, Cook J, Nashelsky J. Vaginal misoprostol for cervical ripening in term pregnancy. *Am.Fam.Physician* 2006;**73**:511-2.

Weeks A, Alfirevic Z. Oral misoprostol administration for labor induction. *Clin.Obstet.Gynecol.* 2006;**49**:658-71.

Weeks A, Alfirevic Z, Faundes A, Hofmeyr GJ, Safar P, Wing D. Misoprostol for induction of labor with a live fetus. *International Journal of Gynaecology & Obstetrics.* **99** Suppl 2:S194-7, 2007 Dec.

Wilson C. The nurse's role in misoprostol induction: a proposed protocol. *J.Obstet.Gynecol.Neonatal Nurs.* 2000;**29**:574-83.

Wing DA. Labor induction with misoprostol. *Am.J.Obstet.Gynecol.* 1999;**181**:339-45.

Wing DA. A benefit-risk assessment of misoprostol for cervical ripening and labour induction. *Drug Saf* 2002;**25**:665-76.

Wing DA, Gaffaney CA. Vaginal misoprostol administration for cervical ripening and labor induction. *Clin. Obstet. Gynecol.* 2006;**49**:627-41.

Winkler M, Rath W. A risk-benefit assessment of oxytocics in obstetric practice. *Drug Safety* 1999;**20**:323-45.

Xenakis EMJ, Piper JM. Chemotherapeutic induction of labour - A rational approach. *Drugs* 1997;**54**:61-8.

Yulzari-Roll V, Hod M, Bar J. [New developments in labor induction]. *Harefuah* 2002;**141**:369-73, 408.

Zust S, Hosli I, Surbek D, Holzgreve W. Side effects of Misoprostol therapy during pregnancy. *Zeitschrift für Geburtshilfe und Neonatologie* 2001;**205**:43-8.

Zust S, Hosli I, Surbek D, Holzgreve W. [Adverse effects of misoprostol in pregnancy]. *Z. Geburtshilfe Neonatol.* 2001;**205**:43-8.

Clinical Studies

Randomised Controlled Trials

Abramovici D, Goldwasser S, Mabie BC, Mercer BM, Goldwasser R, Sibai BM. A randomized comparison of oral misoprostol versus Foley catheter and oxytocin for induction of labor at term. *Am.J. Obstet. Gynecol.* 1999;**181**:1108-12.

Adair CD, Weeks JW, Barrilleaux S, Edwards M, Burlison K, Lewis DF. Oral or vaginal misoprostol administration for induction of labor: a randomized, double-blind trial. *Obstet. Gynecol.* 1998;**92**:810-3.

Adam I, Hassan OA, Elhassan EM. Oral misoprostol vs. vaginal misoprostol for cervical ripening and labor induction. *Int.J. Gynaecol. Obstet.* 2005;**89**:142-3.

Adeniji AO, Olayemi O, Odukogbe AA, Aimakhu CO, Oladokun A, Akindele FO *et al.* Comparison of changes in pre-induction cervical factors' scores following ripening with transcervical foley catheter and intravaginal misoprostol. *Afr.J. Med. Med. Sci.* 2005;**34**:377-82.

Adeniji AO, Olayemi O, Odukogbe AA, Oladokun A, Adeniji OI, Egbewale BE *et al.* Cervico-vaginal foetal fibronectin: a predictor of cervical response at pre-induction cervical ripening. *West Afr. J. Med.* 2005;**24**:334-7.

Adeniji OA, Oladokun A, Olayemi O, Adeniji OI, Odukogbe AA, Ogunbode O *et al.* Pre-induction cervical ripening: transcervical foley catheter versus intravaginal misoprostol. *J.Obstet.Gynaecol.* 2005;**25**:134-9.

Afolabi BB, Oyeneyin OL, Ogedengbe OK. Intravaginal misoprostol versus Foley catheter for cervical ripening and induction of labor. *Int.J.Gynaecol.Obstet.* 2005;**89**:263-7.

Agarwal N, Gupta A, Kriplani A, Bhatla N, Parul. Six hourly vaginal misoprostol versus intracervical dinoprostone for cervical ripening and labor induction. *J.Obstet.Gynaecol.Res.* 2003;**29**:147-51.

Al Hussaini TK, Abdel-Aal SA, Youssef MA. Oral misoprostol vs. intravenous oxytocin for labor induction in women with prelabor rupture of membranes at term. *Int.J.Gynaecol.Obstet.* 2003;**82**:73-5.

Ayad IA. Vaginal misoprostol in managing premature rupture of membranes. *East Mediterr.Health J.* 2002;**8**:515-20.

Barrilleaux PS, Bofill JA, Terrone DA, Magann EF, May WL, Morrison JC. Cervical ripening and induction of labor with misoprostol, dinoprostone gel, and a Foley catheter: a randomized trial of 3 techniques. *Am.J.Obstet.Gynecol.* 2002;**186**:1124-9.

Bartha JL, Comino-Delgado R, Garcia-Benasach F, Martinez-Del-Fresno P, Moreno-Corral LJ. Oral misoprostol and intracervical dinoprostone for cervical ripening and labor induction: a randomized comparison. *Obstet.Gynecol.* 2000;**96**:465-9.

Bartusevicius A, Barcaite E, Krikstolaitis R, Gintautas V, Nadisauskiene R. Sublingual compared with vaginal misoprostol for labour induction at term: a randomised controlled trial. *BJOG.* 2006;**113**:1431-7.

Bebbington M, Schmucl E, Pevzner L, Bernstein P, Dayal A, Barnhard J *et al.* Misoprostol versus dinoprostone for labor induction at term: A randomized controlled trial. *American Journal of Obstetrics and Gynecology* 2003;**189**:S211.

Beigi A, Kabiri M, Zarrinkoub F. Cervical ripening with oral misoprostol at term. *Int.J.Gynaecol.Obstet.* 2003;**83**:251-5.

Belfrage P, Smedvig E, Gjessing L, Eggebo TM, Okland I. A randomized prospective study of misoprostol and dinoprostone for induction of labor. *Acta Obstet.Gynecol.Scand.* 2000;**79**:1065-8.

Bennett KA, Butt K, Crane JM, Hutchens D, Young DC. A masked randomized comparison of oral and vaginal administration of misoprostol for labor induction. *Obstet.Gynecol.* 1998;**92**:481-6.

Bolnick J, Velazquez M, Leslie K, Rappaport R, Gonzalez J, McIlwain G *et al.* Randomized trial of sustained-release vaginal dinoprostone (PGE(2)) with concurrent

oxytocin versus vaginal misoprostol (PGE(1)) for induction of labor term. *Journal of Investigative Medicine* 2003;**51**:S107.

Bolnick JM, Velazquez MD, Gonzalez JL, Rappaport VJ, McIlwain-Dunivan G, Rayburn WF. Randomized trial between two active labor management protocols in the presence of an unfavorable cervix. *Am.J.Obstet.Gynecol.* 2004;**190**:124-8.

Bonebrake R, Haag T, Fleming A, Temp M, Haynatzki G. Vaginal misoprostol is more effective with fewer side effects than oral misoprostol for cervical ripening and induction of labor. *American Journal of Obstetrics and Gynecology* 2001;**185**:S204.

Bower DJ, Meurer LN. Oral versus vaginal administration of misoprostol for labor induction. *J.Fam.Pract.* 1999;**48**:9-10.

Buccellato CA, Stika CS, Frederiksen MC. A randomized trial of misoprostol versus extra-amniotic sodium chloride infusion with oxytocin for induction of labor. *Am.J.Obstet.Gynecol.* 2000;**182**:1039-44.

Buser D, Mora G, Arias F. A randomized comparison between misoprostol and dinoprostone for cervical ripening and labor induction in patients with unfavorable cervixes. *Obstet.Gynecol.* 1997;**89**:581-5.

Butler B, Crane J, Delaney T, Hutchens D, Harnett J. Induction of labour with misoprostol in women at term with an unfavorable cervix: A randomized comparison of oral and vaginal administration. *American Journal of Obstetrics and Gynecology* 2004;**191**:S190.

Butt KD, Bennett KA, Crane JM, Hutchens D, Young DC. Randomized comparison of oral misoprostol and oxytocin for labor induction in term prelabor membrane rupture. *Obstet.Gynecol.* 1999;**94**:994-9.

Byrne JD, Wing DA, Fraser M, Fassett MJ, Goodwin TM, Challis JR. Mifepristone: effect on plasma corticotropin-releasing hormone, adrenocorticotrophic hormone, and cortisol in term pregnancy. *J.Perinatol.* 2004;**24**:416-20.

Caliskan E, Bodur H, Ozeren S, Corakci A, Ozkan S, Yucesoy I. Misoprostol 50 microg sublingually versus vaginally for labor induction at term: a randomized study. *Gynecol.Obstet.Invest* 2005;**59**:155-61.

Campos GA, Guzman S, Rodriguez JG, Voto LS, Margulies M. [Misoprostol--a PGE1 analog for induction of labor at term: comparative and randomized study with oxytocin]. *Rev.Chil.Obstet.Ginecol.* 1994;**59**:190-5.

Carlan SJ, Bouldin S, O'Brien WF. Extemporaneous preparation of misoprostol gel for cervical ripening: a randomized trial. *Obstet.Gynecol.* 1997;**90**:911-5.

Carlan SJ, Bouldin S, Blust D, O'Brien WF. Safety and efficacy of misoprostol orally and vaginally: a randomized trial. *Obstet.Gynecol.* 2001;**98**:107-12.

Carlan SJ, Blust D, O'Brien WF. Buccal versus intravaginal misoprostol administration for cervical ripening. *Am.J.Obstet.Gynecol.* 2002;**186**:229-33.

Cecatti JG, Tedesco RP, Pires HM, Calderon IM, Faundes A. Effectiveness and safety of a new vaginal misoprostol product specifically labeled for cervical ripening and labor induction. *Acta Obstet.Gynecol.Scand.* 2006;**85**:706-11.

Chang CH, Chang FM. Randomized comparison of misoprostol and dinoprostone for preinduction cervical ripening and labor induction. *J.Formos.Med.Assoc.* 1997;**96**:366-9.

Chang YK, Chen WH, Yu MH, Liu HS. Intracervical misoprostol and prostaglandin E2 for labor induction. *Int.J.Gynaecol.Obstet.* 2003;**80**:23-8.

Chanrachakul B, Herabutya Y, Punyavachira P. Randomized trial of isosorbide mononitrate versus misoprostol for cervical ripening at term. *Int.J.Gynaecol.Obstet.* 2002;**78**:139-45.

Chao A, El Sayed YY, Sapugay AM. Does nifedipine prevent the tachysystole associated with misoprostol induction? *J.Matern.Fetal Neonatal Med.* 2002;**11**:266-9.

Charoenkul S, Sripramote M. A randomized comparison of one single dose of vaginal 50 microg misoprostol with 3 mg dinoprostone in pre-induction cervical ripening. *J.Med.Assoc.Thai.* 2000;**83**:1026-34.

Chen DC, Ku CH, Huang YC, Chen CH, Wu GJ. Urinary nitric oxide metabolite changes in spontaneous and induced onset active labor. *Acta Obstetricia et Gynecologica Scandinavica* 2004;**83**:641-6.

Chen DC, Yuan SS, Su HY, Lo SC, Ren SS, Wu GJ. Urinary cyclic guanosine 3',5'-monophosphate and cyclic adenosine 3',5'-monophosphate changes in spontaneous and induced onset active labor. *Acta Obstet.Gynecol.Scand.* 2005;**84**:1081-6.

Cheung PC, Yeo EL, Wong KS, Tang LC. Oral misoprostol for induction of labor in prelabor rupture of membranes (PROM) at term: a randomized control trial. *Acta Obstet.Gynecol.Scand.* 2006;**85**:1128-33.

Chuck FJ, Huffaker BJ. Labor induction with intravaginal misoprostol versus intracervical prostaglandin E2 gel (Prepidil gel): randomized comparison. *Am.J.Obstet.Gynecol.* 1995;**173**:1137-42.

Chung JH, Huang WH, Rumney PJ, Garite TJ, Nageotte MP. A prospective randomized controlled trial that compared misoprostol, Foley catheter, and combination misoprostol-Foley catheter for labor induction. *Am.J.Obstet.Gynecol.* 2003;**189**:1031-5.

- Colon I, Clawson K, Hunter K, Druzin ML, Taslimi MM. Prospective randomized clinical trial of inpatient cervical ripening with stepwise oral misoprostol vs vaginal misoprostol. *Am.J.Obstet.Gynecol.* 2005;**192**:747-52.
- Crane JM, Delaney T, Hutchens D. Oral misoprostol for premature rupture of membranes at term. *Am.J.Obstet.Gynecol.* 2003;**189**:720-4.
- Culver J, Strauss RA, Brody S, Dorman K, Timlin S, McMahon MJ. A randomized trial comparing vaginal misoprostol versus Foley catheter with concurrent oxytocin for labor induction in nulliparous women. *Am.J.Perinatol.* 2004;**21**:139-46.
- da Graca KF, Cecatti JG, Castro Surita FG, Milanez HM, Parpinelli MA. Misoprostol versus expectant management in premature rupture of membranes at term. *BJOG.* 2005;**112**:1284-90.
- Dallen-Bach P, Boulvain M, Viardot C, Irion O. Oral Misoprostol or vaginal Dinoprostone for labor induction? Randomized controlled trial. *American Journal of Obstetrics and Gynecology* 2001;**185**:S108.
- Danielian P, Porter B, Ferri N, Summers J, Templeton A. Misoprostol for induction of labour at term: a more effective agent than dinoprostone vaginal gel. *Br.J.Obstet.Gynaecol.* 1999;**106**:793-7.
- de Aquino MM, Cecatti JG. Misoprostol versus oxytocin for labor induction in term and post-term pregnancy: randomized controlled trial. *Sao Paulo Med.J.* 2003;**121**:102-6.
- de la TS, Gilson GJ, Flores S, Curet LB, Qualls CE, Rayburn WF. Is high-dose misoprostol able to lower the incidence of cesarean section? A randomized controlled trial. *J.Matern.Fetal Med.* 2001;**10**:85-90.
- De A, Bagga R, Gopalan S. The routine use of oxytocin after oral misoprostol for labour induction in women with an unfavourable cervix is not of benefit. *Aust.N.Z.J.Obstet.Gynaecol.* 2006;**46**:323-9.
- Delaney T, Crane JM, Hutchens D, Fanning CA, Young DC. Induction of labor with intravaginal misoprostol: A comparison of dosing intervals. *American Journal of Obstetrics and Gynecology* 2001;**185**:S202.
- Denguezli W, Trimech A, Haddad A, Hajjaji A, Saidani Z, Faleh R *et al.* Efficacy and safety of six hourly vaginal misoprostol versus intracervical dinoprostone: a randomized controlled trial. *Arch.Gynecol.Obstet.* 2007.
- Diro M, Adra A, Gilles JM, Nassar A, Rodriguez A, Salamat SM *et al.* A double-blind randomized trial of two dose regimens of misoprostol for cervical ripening and labor induction. *J.Matern.Fetal Med.* 1999;**8**:114-8.

Dodd JM, Crowther CA, Robinson JS. Morning compared with evening induction of labor: a nested randomized controlled trial. A nested randomized controlled trial. *Obstet.Gynecol.* 2006;**108**:350-60.

Dodd JM, Crowther CA, Robinson JS. Oral misoprostol for induction of labour at term: randomised controlled trial. *British Medical Journal* 2006;**332**:509-11.

Dodd JM, Crowther CA, Robinson JS. Oral misoprostol for induction of labour at term: randomised controlled trial. *BMJ* 2006;**332**:509-13.

Echeverria E, Rocha M. [Randomized comparative study of induced labor with oxytocin and misoprostol in prolonged pregnancies]. *Rev.Chil.Obstet.Ginecol.* 1995;**60**:108-11.

El Sherbiny MT, El Gharieb IH, Gewely HA. Vaginal misoprostol for induction of labor: 25 vs. 50 microg dose regimen. *Int.J.Gynaecol.Obstet.* 2001;**72**:25-30.

Elhassan EM, Mirghani OA, Adam I. Misoprostol vs. oxytocin for induction of labor. *Int.J.Gynaecol.Obstet.* 2005;**91**:254-5.

Elhassan EM, Mirghani OA, Adam I. Cervical ripening and labor induction with 25 microg vs. 50 microg of intravaginal misoprostol. *Int.J.Gynaecol.Obstet.* 2005;**90**:234-5.

Elhassan EM, Nasr AM, Adam I. Sublingual compared with oral and vaginal misoprostol for labor induction. *Int.J.Gynaecol.Obstet.* 2007;**97**:153-4.

Elhassan M, Mirghani OA, Adam I. Intravaginal misoprostol vs. dinoprostone as cervical ripening and labor-inducing agents. *Int.J.Gynaecol.Obstet.* 2004;**85**:285-6.

Elhassan EM, Nasr AM, Adam I. Sublingual compared with oral and vaginal misoprostol for labor induction. *International Journal of Gynaecology & Obstetrics.* 97(2):153-4, 2007 May.

Eroglu D, Oktem M, Yanik F, Kuscu E. Labor induction at term: a comparison of the effects of 50 microg and 25 microg vaginal misoprostol. *Clinical & Experimental Obstetrics & Gynecology.* 34(2):102-5, 2007.

Escudero F, Contreras H. A comparative trial of labor induction with misoprostol versus oxytocin. *Int.J.Gynaecol.Obstet.* 1997;**57**:139-43.

Ewert K, Powers B, Robertson S, Alfirevic Z. Controlled-release misoprostol vaginal insert in parous women for labor induction: a randomized controlled trial. *Obstet.Gynecol.* 2006;**108**:1130-7.

- Farah LA, Sanchez-Ramos L, Rosa C, Del Valle GO, Gaudier FL, Delke I *et al.* Randomized trial of two doses of the prostaglandin E1 analog misoprostol for labor induction. *Am.J.Obstet.Gynecol.* 1997;**177**:364-9.
- Feitosa FE, Sampaio ZS, Alencar CA, Jr., Amorim MM, Passini R, Jr. Sublingual vs. vaginal misoprostol for induction of labor. *Int.J.Gynaecol.Obstet.* 2006;**94**:91-5.
- Ferguson JE, Head BH, Frank FH, Frank ML, Singer JS, Stefos T *et al.* Misoprostol versus low-dose oxytocin for cervical ripening: a prospective, randomized, double-masked trial. *Am.J.Obstet.Gynecol.* 2002;**187**:273-9.
- Fischer J, Newton W. Should we use Foley catheters for preinduction cervical ripening? *J.Fam.Pract.* 2001;**50**:655.
- Fisher SA, Mackenzie VP, Davies GA. Oral versus vaginal misoprostol for induction of labor: a double-blind randomized controlled trial. *Am.J.Obstet.Gynecol.* 2001;**185**:906-10.
- Fletcher H, Mitchell S, Frederick J, Simeon D, Brown D. Intravaginal misoprostol versus dinoprostone as cervical ripening and labor-inducing agents. *Obstet.Gynecol.* 1994;**83**:244-7.
- Fletcher HM, Mitchell S, Simeon D, Frederick J, Brown D. Intravaginal misoprostol as a cervical ripening agent. *Br.J.Obstet.Gynaecol.* 1993;**100**:641-4.
- Frohn WE, Simmons S, Carlan SJ. Prostaglandin E-2 gel versus misoprostol for cervical ripening in patients with premature rupture of membranes after 34 weeks. *Obstetrics and Gynecology* 2002;**99**:206-10.
- Frohn WE, Simmons S, Carlan SJ. Prostaglandin E2 gel versus misoprostol for cervical ripening in patients with premature rupture of membranes after 34 weeks. *Obstet.Gynecol.* 2002;**99**:206-10.
- Garry D, Figueroa R, Kalish RB, Catalano CJ, Maulik D. Randomized controlled trial of vaginal misoprostol versus dinoprostone vaginal insert for labor induction. *J.Matern.Fetal Neonatal Med.* 2003;**13**:254-9.
- Gelisen O, Caliskan E, Dilbaz S, Ozdas E, Dilbaz B, Ozdas E *et al.* Induction of labor with three different techniques at 41 weeks of gestation or spontaneous follow-up until 42 weeks in women with definitely unfavorable cervical scores. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2005;**120**:164-9.
- Gherman RB, Browning J, O'Boyle A, Goodwin TM. Oral misoprostol vs. intravaginal prostaglandin E2 for preinduction cervical ripening. A randomized trial. *J.Reprod.Med.* 2001;**46**:641-6.

- Ghidini A, Spong CY, Korker V, Mariani E. Randomized controlled trial of 50 and 100 mcg of misoprostol for induction of labor at term. *Arch.Gynecol.Obstet.* 2001;**265**:128-30.
- Gilson GJ, Russell DJ, Izquierdo LA, Qualls CR, Curet LB. A prospective randomized evaluation of a hygroscopic cervical dilator, Dilapan, in the preinduction ripening of patients undergoing induction of labor. *American Journal of Obstetrics and Gynecology* 1996;**175**:145-9.
- Gottschall DS, Borgida AF, Mihalek JJ, Sauer F, Rodis JF. A randomized clinical trial comparing misoprostol with prostaglandin E2 gel for preinduction cervical ripening. *Am.J.Obstet.Gynecol.* 1997;**177**:1067-70.
- Gregson S, Waterstone M, Norman I, Murrells T. A randomised controlled trial comparing low dose vaginal misoprostol and dinoprostone vaginal gel for inducing labour at term. *BJOG.* 2005;**112**:438-44.
- Gregson S, Waterstone M, Norman I, Murrells T. A randomised controlled trial comparing low dose vaginal misoprostol and dinoprostone vaginal gel for inducing labour at term. *Bjog-An International Journal of Obstetrics and Gynaecology* 2005;**112**:438-44.
- Greybush M, Singleton C, Atlas RO, Balducci J, Rust OA. Preinduction cervical ripening techniques compared. *J.Reprod.Med.* 2001;**46**:11-7.
- Haghighi L. Intravaginal misoprostol in preterm premature rupture of membranes with low Bishop scores. *Int.J.Gynaecol.Obstet.* 2006;**94**:121-2.
- Hall R, Duarte-Gardea M, Harlass F. Oral versus vaginal misoprostol for labor induction. *Obstet.Gynecol.* 2002;**99**:1044-8.
- Has R, Batukan C, Ermis H, Cevher E, Araman A, Kilic G *et al.* Comparison of 25 and 50 microg vaginally administered misoprostol for preinduction of cervical ripening and labor induction. *Gynecol.Obstet.Invest* 2002;**53**:16-21.
- Herabutya Y, Prasertsawat P, Pokpirom J. A comparison of intravaginal misoprostol and intracervical prostaglandin E2 gel for ripening of unfavorable cervix and labor induction. *J.Obstet.Gynaecol.Res.* 1997;**23**:369-74.
- Hoffmann RA, Anthony J, Fawcus S. Oral misoprostol vs. placebo in the management of prelabor rupture of membranes at term. *Int.J.Gynaecol.Obstet.* 2001;**72**:215-21.
- Hofmeyr GJ, Alfrevic Z, Matonhodze B, Brocklehurst P, Campbell E, Nikodem VC. Titrated oral misoprostol solution for induction of labour: a multi-centre, randomised trial. *BJOG.* 2001;**108**:952-9.

How HY, Leaseburge L, Khoury JC, Siddiqi TA, Spinnato JA, Sibai BM. A comparison of various routes and dosages of misoprostol for cervical ripening and the induction of labor. *Am.J.Obstet.Gynecol.* 2001;**185**:911-5.

Incerpi MH, Fassett MJ, Kjos SL, Tran SH, Wing DA. Vaginally administered misoprostol for outpatient cervical ripening in pregnancies complicated by diabetes mellitus. *Am.J.Obstet.Gynecol.* 2001;**185**:916-9.

Jazayeri A, Jazayeri M, Jamal A, Eslamian L, Marsoosi V, Borna S. Prospective randomized clinical trial of cervical ripening with misoprostol for either 8 or 24 hours. *American Journal of Obstetrics and Gynecology* 2003;**189**:S70.

Jiang X, Wang H, Zhang Z. [Determination of fetal umbilical artery flow velocity during induction of term labor by mifepristone]. *Zhonghua Fu Chan Ke.Za Zhi.* 1997;**32**:732-4.

Kadanali S, Kucukozkan T, Zor N, Kumtepe Y. Comparison of labor induction with misoprostol vs. oxytocin/prostaglandin E2 in term pregnancy. *Int.J.Gynaecol.Obstet.* 1996;**55**:99-104.

Karjane NW, Brock EL, Walsh SW. Induction of labor using a Foley balloon, with and without extra-amniotic saline infusion. *Obstetrics and Gynecology* 2006;**107**:234-9.

Kashanian M, Akbarian AR, Fekrat M. Cervical ripening and induction of labor with intravaginal misoprostol and Foley catheter cervical traction. *Int.J.Gynaecol.Obstet.* 2006;**92**:79-80.

Khoury AN, Zhou QP, Gorenberg DM, Nies BM, Manley GE, Mecklenburg FE. A comparison of intermittent vaginal administration of two different doses of misoprostol suppositories with continuous dinoprostone for cervical ripening and labor induction. *J.Matern.Fetal Med.* 2001;**10**:186-92.

Kidanto HL, Kaguta MM, van Roosmalen J. Induction of labor with misoprostol or oxytocin in Tanzania. *Int.J.Gynaecol.Obstet.* 2007;**96**:30-1.

Kipikasa JH, Adair CD, Williamson J, Breen JM, Medford LK, Sanchez-Ramos L. Use of misoprostol on an outpatient basis for postdate pregnancy. *Int.J.Gynaecol.Obstet.* 2005;**88**:108-11.

Kolderup L, McLean L, Grullon K, Safford K, Kilpatrick SJ. Misoprostol is more efficacious for labor induction than prostaglandin E2, but is it associated with more risk? *Am.J.Obstet.Gynecol.* 1999;**180**:1543-50.

Kramer RL, Gilson GJ, Morrison DS, Martin D, Gonzales JL, Qualls CR. A randomized trial of misoprostol and oxytocin for induction of labor: safety and efficacy. *Obstet.Gynecol.* 1997;**89**:387-91.

Kulshreshtha S, Sharma P, Mohan G, Singh S, Singh S. Comparative study of misoprostol vs dinoprostone for induction of labour. *Indian Journal of Physiology & Pharmacology*. 51(1):55-61, 2007 Jan-Mar.

Kwon JS, Davies GA, Mackenzie VP. A comparison of oral and vaginal misoprostol for induction of labour at term: a randomised trial. *BJOG*. 2001;108:23-6.

Langenegger EJ, Odendaal HJ, Grove D. Oral misoprostol versus intracervical dinoprostone for induction of labor. *Int.J.Gynaecol.Obstet*. 2005;88:242-8.

Lapaire O, Zanetti-Dallenbach R, Weber P, Hosli I, Holzgreve W, Surbek D. Labor induction in preeclampsia: is misoprostol more effective than dinoprostone?. *Journal of Perinatal Medicine*. 35(3):195-9, 2007

Laube DW. Induction of labor. *Clinical Obstetrics and Gynecology* 1997;40:485-95.

le Roux PA, Olarogun JO, Penny J, Anthony J. Oral and vaginal misoprostol compared with dinoprostone for induction of labor: a randomized controlled trial. *Obstet.Gynecol*. 2002;99:201-5.

Lee HY. A randomised double-blind study of vaginal misoprostol vs dinoprostone for cervical ripening and labour induction in prolonged pregnancy. *Singapore Med.J*. 1997;38:292-4.

Lemancewicz A, Urban R, Skotnicki MZ, Karpiuk A, Urban J. Uterine and fetal Doppler flow changes after misoprostol and oxytocin therapy for induction of labor in post-term pregnancies. *Int.J.Gynaecol.Obstet*. 1999;67:139-45.

Leszczynska-Gorzela B, Laskowska M, Oleszczuk J. [Using of Misoprostol for preinduction and induction of labor in term pregnancy]. *Ginekol.Pol*. 1999;70:881-9.

Leszczynska-Gorzela B, Laskowska M, Oleszczuk J. Comparative analysis of the effectiveness of misoprostol and prostaglandin E(2) in the preinduction and induction of labor. *Med.Sci.Monit*. 2001;7:1023-8.

Levy R, Kanengiser B, Furman B, Ben Arie A, Brown D, Hagay ZJ. A randomized trial comparing a 30-mL and an 80-mL Foley catheter balloon for preinduction cervical ripening. *American Journal of Obstetrics and Gynecology* 2004;191:1632-6.

Levy R, Vaisbuch E, Furman B, Doitch H, Oron S, Hagay Z. Prospective randomized clinical trial of immediate induction of labor with oral misoprostol for prelabor rupture of the membranes in women with unfavorable cervix at term. *American Journal of Obstetrics and Gynecology* 2005;193:S44.

Levy R, Vaisbuch E, Furman B, Brown D, Volach V, Hagay ZJ. Induction of labor with oral misoprostol for premature rupture of membranes at term in women with

unfavorable cervix: a randomized, double-blind, placebo-controlled trial. *J.Perinat.Med.* 2007;**35**:126-9.

Lo JY, Alexander JM, McIntire DD, Leveno KJ. Efficacy of oral misoprostol in nulliparous women with premature rupture of membranes. *American Journal of Obstetrics and Gynecology* 2001;**185**:S204.

Lo JY, Alexander JM, McIntire DD, Leveno KJ. Randomized trial of oral misoprostol in nulliparous women with premature rupture of membranes at term. *American Journal of Obstetrics and Gynecology* 2001;**185**:S204.

Lo JY, Alexander JM, McIntire DD, Leveno KJ. Ruptured membranes at term: randomized, double-blind trial of oral misoprostol for labor induction. *Obstet.Gynecol.* 2003;**101**:685-9.

Lo TK, Lau WL, Wong KS, Tang LC. Sublingual misoprostol compared to artificial rupture of membranes plus oxytocin infusion for labour induction in nulliparous women with a favourable cervix at term. *Hong.Kong.Med.J.* 2006;**12**:345-50.

Lokugamage AU, Forsyth SF, Sullivan KR, El Refaey H, Rodeck CH. Dinoprostone versus misoprostol: a randomized study of nulliparous women undergoing induction of labor. *Acta Obstet.Gynecol.Scand.* 2003;**82**:133-7.

Lokugamage AU, Forsyth SF, Sullivan KR, El Refaey H, Rodeck CH. Randomized trial in multiparous patients: investigating a single vs. two-dose regimen of intravaginal misoprostol for induction of labor. *Acta Obstet.Gynecol.Scand.* 2003;**82**:138-42.

Magtibay PM, Ramin KD, Harris DY, Ramsey PS, Ogburn PL, Jr. Misoprostol as a labor induction agent. *J.Matern.Fetal Med.* 1998;**7**:15-8.

Majoko F, Nystrom L, Lindmark G. No benefit, but increased harm from high dose (100 microg) misoprostol for induction of labour: a randomised trial of high vs. low (50 microg) dose misoprostol. *J.Obstet.Gynaecol.* 2002;**22**:614-7.

Majoko F, Zwizwai M, Lindmark G, Nystrom L. Labor induction with vaginal misoprostol and extra-amniotic prostaglandin F2alpha gel. *Int.J.Gynaecol.Obstet.* 2002;**76**:127-33.

Majoko F, Zwizwai M, Nystrom L, Lindmark G. Vaginal misoprostol for induction of labour: a more effective agent than prostaglandin F2 alpha gel and prostaglandin E2 pessary. *Cent.Afr.J.Med.* 2002;**48**:123-8.

Margulies M, Campos PG, Voto LS. Misoprostol to induce labour. *Lancet* 1992;**339**:64.

- Matonhodze BB, Hofmeyr GJ, Levin J. Labour induction at term--a randomised trial comparing Foley catheter plus titrated oral misoprostol solution, titrated oral misoprostol solution alone, and dinoprostone. *S.Afr.Med.J.* 2003;**93**:375-9.
- McKenna DS, Ester JB, Proffitt M, Waddell KR. Misoprostol outpatient cervical ripening without subsequent induction of labor: a randomized trial. *Obstet.Gynecol.* 2004;**104**:579-84.
- Megalo A, Petignat P, Hohlfeld P. Influence of misoprostol or prostaglandin E(2) for induction of labor on the incidence of pathological CTG tracing: a randomized trial. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2004;**116**:34-8.
- Meydanli MM, Caliskan E, Burak F, Narin MA, Atmaca R. Labor induction post-term with 25 micrograms vs. 50 micrograms of intravaginal misoprostol. *Int.J.Gynaecol.Obstet.* 2003;**81**:249-55.
- Meyer M, Pflum J, Howard D. Outpatient misoprostol compared with dinoprostone gel for preinduction cervical ripening: a randomized controlled trial. *Obstet.Gynecol.* 2005;**105**:466-72.
- Milchev N, Kuzmanov B, Terzhumanov R. [Cytotec: an effective drug for the induction of labor]. *Akush.Ginekol.(Sofia)* 2003;**42**:9-11.
- Moodley J, Venkatachalam S, Songca P. Misoprostol for cervical ripening at and near term--a comparative study. *S.Afr.Med.J.* 2003;**93**:371-4.
- Morgan OF, Baez BJ, Quevedo CE, Cuetos Martinez CB, Osuna R, I. [Misoprostol and oxytocin for induction of cervical ripening and labor in patients with term pregnancy and premature membrane rupture]. *Ginecol.Obstet.Mex.* 2002;**70**:469-76.
- Mozurkewich E, Horrocks J, Daley S, Von Oeyen P, Halvorson M, Johnson M *et al.* The MisoPROM study: a multicenter randomized comparison of oral misoprostol and oxytocin for premature rupture of membranes at term. *Am.J.Obstet.Gynecol.* 2003;**189**:1026-30.
- Mullin PM, House M, Paul RH, Wing DA. A comparison of vaginally administered misoprostol with extra-amniotic saline solution infusion for cervical ripening and labor induction. *Am.J.Obstet.Gynecol.* 2002;**187**:847-52.
- Mundle WR, Young DC. Vaginal misoprostol for induction of labor: a randomized controlled trial. *Obstet.Gynecol.* 1996;**88**:521-5.
- Nanda S, Singhal SR, Papneja A. Induction of labour with intravaginal misoprostol and prostaglandin E2 gel: a comparative study. *Trop.Doct.* 2007;**37**:21-4.
- Nassar AH, Awwad J, Khalil AM, Abu-Musa A, Mehio G, Usta IM. A randomised comparison of patient satisfaction with vaginal and sublingual misoprostol for

induction of labour at term. *BJOG: An International Journal of Obstetrics & Gynaecology*. **114**(10):1215-21, 2007 Oct.

Neiger R, Greaves PC. Comparison between vaginal misoprostol and cervical dinoprostone for cervical ripening and labor induction. *Tenn.Med.* 2001;**94**:25-7.

Ngai SW, To WK, Lao T, Ho PC. Cervical priming with oral misoprostol in pre-labor rupture of membranes at term. *Obstet.Gynecol.* 1996;**87**:923-6.

Ngai SW, Chan YM, Lam SW, Lao TT. Labour characteristics and uterine activity: misoprostol compared with oxytocin in women at term with prelabour rupture of the membranes. *BJOG*. 2000;**107**:222-7.

Nigam A, Singh VK, Dubay P, Pandey K, Bhagoliwal A, Prakash A. Misoprostol vs. oxytocin for induction of labor at term. *Int.J.Gynaecol.Obstet.* 2004;**86**:398-400.

Nopdonrattakoon L. A comparison between intravaginal and oral misoprostol for labor induction: a randomized controlled trial. *J.Obstet.Gynaecol.Res.* 2003;**29**:87-91.

Nunes F, Rodrigues R, Meirinho M. Randomized comparison between intravaginal misoprostol and dinoprostone for cervical ripening and induction of labor. *Am.J.Obstet.Gynecol.* 1999;**181**:626-9.

Nuthalapaty F, Ramsey P, Biggio J, Owen J. Comparative efficacy of high dose vaginal misoprostol versus concentrated oxytocin plus low dose vaginal misoprostol for midtrimester labor induction. *American Journal of Obstetrics and Gynecology* 2004;**191**:S73.

O'Flaherty L. Oral misoprostol and intracervical dinoprostone for cervical ripening and labor induction: a randomized comparison. *J.Midwifery Womens Health* 2001;**46**:203-4.

O'Sullivan MJ, Frank F. Misoprostol versus low-dose oxytocin for cervical ripening: A prospective, randomized, double-masked trial - Discussion. *American Journal of Obstetrics and Gynecology* 2002;**187**:279-80.

Oboro VO, Tabowei TO. Outpatient misoprostol cervical ripening without subsequent induction of labor to prevent post-term pregnancy. *Acta Obstet.Gynecol.Scand.* 2005;**84**:628-31.

Owolabi AT, Kuti O, Ogunlola IO. Randomised trial of intravaginal misoprostol and intracervical Foley catheter for cervical ripening and induction of labour. *J.Obstet.Gynaecol.* 2005;**25**:565-8.

Ozgun K, Kizilates A, Uner M, Erman O, Trak B. Induction of labor with intravaginal misoprostol versus intracervical dinoprostone. *Arch.Gynecol.Obstet.* 1997;**261**:9-13.

- Ozsoy M, Ozsoy D. Induction of labor with 50 and 100 microg of misoprostol: comparison of maternal and fetal outcomes. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2004;**113**:41-4.
- Pandis GK, Papageorghiou AT, Otigbah CM, Howard RJ, Nicolaides KH. Randomized study of vaginal misoprostol (PGE(1)) and dinoprostone gel (PGE(2)) for induction of labor at term. *Ultrasound Obstet.Gynecol.* 2001;**18**:629-35.
- Papanikolaou EG, Plachouras N, Drougia A, Andronikou S, Vlachou C, Stefos T *et al.* Comparison of misoprostol and dinoprostone for elective induction of labour in nulliparous women at full term: a randomized prospective study. *Reprod.Biol.Endocrinol.* 2004;**2**:70.
- Paungmora N, Herabutya Y, Prasertsawat P, Punyavachira P. Comparison of oral and vaginal misoprostol for induction of labor at term: a randomized controlled trial. *J.Obstet.Gynaecol.Res.* 2004;**30**:358-62.
- Perry KG, Jr., Larmon JE, May WL, Robinette LG, Martin RW. Cervical ripening: a randomized comparison between intravaginal misoprostol and an intracervical balloon catheter combined with intravaginal dinoprostone. *Am.J.Obstet.Gynecol.* 1998;**178**:1333-40.
- Pi P, Zhu F. [Clinical observation of misoprostol on induction in late pregnancy]. *Hunan.Yi.Ke.Da.Xue.Xue.Bao.* 1999;**24**:195-7.
- Pongsatha S, Tongsong T, Somsak T. A comparison between 50 mcg oral misoprostol every 4 hours and 6 hours for labor induction: a prospective randomized controlled trial. *J.Med.Assoc.Thai.* 2001;**84**:989-94.
- Pongsatha S, Sirisukkasem S, Tongsong T. A comparison of 100 microg oral misoprostol every 3 hours and 6 hours for labor induction: a randomized controlled trial. *J.Obstet.Gynaecol.Res.* 2002;**28**:308-12.
- Pongsatha S, Vijittrawiwat A, Tongsong T. A comparison of labor induction by oral and vaginal misoprostol. *Int.J.Gynaecol.Obstet.* 2005;**88**:140-1.
- Porozhanova V, Sampat D, Porozhanova K. [Misoprostol and induction of labour]. *Akush.Ginekol.(Sofia)* 2005;**44**:27-30.
- Raio L, Ghezzi F, Di Naro E, Bruhwiler H, Luscher KP. [Shorter delivery time after induction with misoprostol]. *Z.Geburtshilfe Neonatol.* 2001;**205**:147-51.
- Ramsey P, Savage K, Owen J. Comparative efficacy of high-dose vaginal misoprostol versus concentrated oxytocin plus low-dose vaginal PGE2 for mid-trimester labor induction. *American Journal of Obstetrics and Gynecology* 2001;**185**:S202.

Ramsey PS, Harris DY, Ogburn PL, Heise RH, Magtibay PM, Ramin KD. Comparative cost analysis of prostaglandin analogues dinoprostone and misoprostol as labor preinduction agents. *Prim.Care Update.Ob Gyns.* 1998;**5**:182.

Ramsey PS, Harris DY, Ogburn PL, Jr., Heise RH, Magtibay PM, Ramin KD. Comparative efficacy and cost of the prostaglandin analogs dinoprostone and misoprostol as labor preinduction agents. *Am.J.Obstet.Gynecol.* 2003;**188**:560-5.

Ramsey PS, Meyer L, Walkes BA, Harris D, Ogburn PL, Jr., Heise RH *et al.* Cardiotocographic abnormalities associated with dinoprostone and misoprostol cervical ripening. *Obstet.Gynecol.* 2005;**105**:85-90.

Rasheed R. Alam AA. Younus S. Raza F. Oral versus vaginal misoprostol for labour induction. *JPMA - Journal of the Pakistan Medical Association.* **57**(8):404-7, 2007 Aug.

Reyna-Villasmil E, Guerra-Velasquez M, Torres-Montilla M, Reyna-Villasmil N, Mejia-Montilla J, Labarca-Vincero N. [Comparative study of the effect of intravaginal misoprostol at 50 and 100 micrograms in cervical ripening and labor induction]. *Invest Clin.* 2005;**46**:179-86.

Rockhead C, Fletcher H, Reid M, Morgan O. A comparison of two methods of labor induction with vaginal misoprostol. *Int.J.Gynaecol.Obstet.* 2003;**80**:271-7.

Rowlands S, Bell R, Donath S, Morrow S, Trudinger BJ. Misoprostol versus dinoprostone for cervical priming prior to induction of labour in term pregnancy: a randomised controlled trial. *Aust.N.Z.J.Obstet.Gynaecol.* 2001;**41**:145-52.

Rowles RB, Paul RH, Nichols MD, Goodwin TM, Schroeder PW, Parer JT *et al.* Misoprostol is more efficacious for labor induction than prostaglandin E-2, but is it associated with more risk? Discussion. *American Journal of Obstetrics and Gynecology* 1999;**180**:1548-50.

Rozenberg P, Chevret S, Goffinet F, Durand-Zaleski I, Ville Y, Vayssiere C *et al.* Induction of labour with a viable infant: a randomised clinical trial comparing intravaginal misoprostol and intravaginal dinoprostone. *BJOG.* 2001;**108**:1255-62.

Rozenberg P, Chevret S, Senat MV, Bretelle F, Paule BA, Ville Y. A randomized trial that compared intravaginal misoprostol and dinoprostone vaginal insert in pregnancies at high risk of fetal distress. *Am.J.Obstet.Gynecol.* 2004;**191**:247-53.

Rust OA, Greybush M, Atlas RO, Jones KJ, Balducci J. Preinduction cervical ripening. A randomized trial of intravaginal misoprostol alone vs. a combination of transcervical Foley balloon and intravaginal misoprostol. *J.Reprod.Med.* 2001;**46**:899-904.

Sahin HG, Sahin HA, Kocer M. Induction of labor in toxemia with misoprostol. *Acta Obstet.Gynecol.Scand.* 2002;**81**:252-7.

Saleem S. Efficacy of dinoprostone, intracervical foleys and misoprostol in labor induction. *J.Coll.Physicians Surg.Pak.* 2006;**16**:276-9.

Sanchez-Ramos L, Kaunitz AM, Del Valle GO, Delke I, Schroeder PA, Briones DK. Labor induction with the prostaglandin E1 methyl analogue misoprostol versus oxytocin: a randomized trial. *Obstet.Gynecol.* 1993;**81**:332-6.

Sanchez-Ramos L, Chen AH, Kaunitz AM, Gaudier FL, Delke I. Labor induction with intravaginal misoprostol in term premature rupture of membranes: a randomized study. *Obstet.Gynecol.* 1997;**89**:909-12.

Sanchez-Ramos L, Peterson DE, Delke I, Gaudier FL, Kaunitz AM. Labor induction with prostaglandin E1 misoprostol compared with dinoprostone vaginal insert: a randomized trial. *Obstet.Gynecol.* 1998;**91**:401-5.

Sanchez-Ramos L, Danner CJ, Delke I, Kaunitz AM. The effect of tablet moistening on labor induction with intravaginal misoprostol: a randomized trial. *Obstet.Gynecol.* 2002;**99**:1080-4.

Sanchez-Ramos L. Induction of labor. *Obstetrics and Gynecology Clinics of North America* 2005;**32**:181-+.

Schaff EA, DiCenzo R, Fielding SL. Comparison of misoprostol plasma concentrations following buccal and sublingual administration. *Contraception* 2005;**71**:22-5.

Schneider M, Ramsey R, Kao L, Bennett KA. Misoprostol is effective for induction of labor in high risk pregnant women: A randomized controlled trial. *American Journal of Obstetrics and Gynecology* 2004;**191**:S73.

Sciscione AC, Nguyen L, Manley J, Pollock M, Maas B, Colmorgen G. A randomized comparison of transcervical Foley catheter to intravaginal misoprostol for preinduction cervical ripening. *Obstet.Gynecol.* 2001;**97**:603-7.

Sharami SH, Milani F, Zahiri Z, Mansour-Ghanaei F. A randomized trial of prostaglandin E-2 gel and extra-amniotic saline infusion with high dose oxytocin for cervical ripening. *Medical Science Monitor* 2005;**11**:CR381-CR386.

Sharma Y, Kumar S, Mittal S, Misra R, Dadhwal V. Evaluation of glyceryl trinitrate, misoprostol, and prostaglandin E2 gel for preinduction cervical ripening in term pregnancy. *J.Obstet.Gynaecol.Res.* 2005;**31**:210-5.

Sherman DJ, Frenkel E, Pansky M, Caspi E, Bukovsky I, Langer R. Balloon cervical ripening with extra-amniotic infusion of saline or prostaglandin E-2: A double-blind, randomized controlled study. *Obstetrics and Gynecology* 2001;**97**:375-80.

- Shetty A, Danielian P, Templeton A. Oral vs vaginal misoprostol in the induction of labour at term: a randomised controlled trial. *British Journal of Obstetrics and Gynaecology* 2000;**107**:813.
- Shetty A, Danielian P, Templeton A. A comparison of oral and vaginal misoprostol tablets in induction of labour at term. *BJOG*. 2001;**108**:238-43.
- Shetty A, Martin R, Danielian P, Templeton A. A comparison of two dosage regimens of oral misoprostol for labor induction at term. *Acta Obstet.Gynecol.Scand*. 2002;**81**:337-42.
- Shetty A, Danielian P, Templeton A. Sublingual misoprostol for the induction of labor at term. *Am.J.Obstet.Gynecol*. 2002;**186**:72-6.
- Shetty A, Stewart K, Stewart G, Rice P, Danielian P, Templeton A. Active management of term prelabour rupture of membranes with oral misoprostol. *BJOG*. 2002;**109**:1354-8.
- Shetty A, Mackie L, Danielian P, Rice P, Templeton A. Sublingual compared with oral misoprostol in term labour induction: a randomised controlled trial. *BJOG*. 2002;**109**:645-50.
- Shetty A, Livingstone I, Acharya S, Rice P, Danielian P, Templeton A. Oral misoprostol (100 microg) versus vaginal misoprostol (25 microg) in term labor induction: a randomized comparison. *Acta Obstet.Gynecol.Scand*. 2003;**82**:1103-6.
- Shetty A, Livingstone I, Acharya S, Rice P, Danielian P, Templeton A. A randomised comparison of oral misoprostol and vaginal prostaglandin E2 tablets in labour induction at term. *BJOG*. 2004;**111**:436-40.
- Sifakis S, Angelakis E, Avgoustinakis E, Fragouli Y, Mantas N, Koukoura O *et al*. A randomized comparison between intravaginal misoprostol and prostaglandin E(2) for labor induction. *Arch.Gynecol.Obstet*. 2007;**275**:263-7.
- Srisomboon J, Tongsong T, Tosiri V. Preinduction cervical ripening with intravaginal prostaglandin E1 methyl analogue misoprostol: a randomized controlled trial. *J.Obstet.Gynaecol.Res*. 1996;**22**:119-24.
- Srisomboon J, Piyamongkol W, Aiewsakul P. Comparison of intracervical and intravaginal misoprostol for cervical ripening and labour induction in patients with an unfavourable cervix. *J.Med.Assoc.Thai*. 1997;**80**:189-94.
- Srisomboon J, Singchai S. A comparison between 25 micrograms and 50 micrograms of intravaginal misoprostol for labor induction. *J.Med.Assoc.Thai*. 1998;**81**:779-83.
- Stitely ML, Browning J, Fowler M, Gendron RT, Gherman RB. Outpatient cervical ripening with intravaginal misoprostol. *Obstet.Gynecol*. 2000;**96**:684-8.

Surbek DV, Boesiger H, Hoesli I, Pavic N, Holzgreve W. A double-blind comparison of the safety and efficacy of intravaginal misoprostol and prostaglandin E2 to induce labor. *Am.J.Obstet.Gynecol.* 1997;**177**:1018-23.

Tabowei TO, Oboro VO. Low dose intravaginal misoprostol versus intracervical balloon catheter for pre-induction cervical ripening. *East Afr.Med.J.* 2003;**80**:91-4.

Thigpen B, Bofill J, Bufkin L, Woodring T, Moore L, Morrison J. A randomized clinical trial comparing vaginal misoprostol to cervical Foley plus oral misoprostol for cervical ripening and labor induction. *American Journal of Obstetrics and Gynecology* 2004;**191**:S18.

Topozada MK, Anwar MY, Hassan HA, el Gazerly WS. Oral or vaginal misoprostol for induction of labor. *Int.J.Gynaecol.Obstet.* 1997;**56**:135-9.

Tukur J, Umar NI, Khan N, Musa D. Comparison of emergency caesarean section to misoprostol induction for the delivery of antepartum eclamptic patients: a pilot study. *Nigerian Journal of Medicine: Journal of the National Association of Resident Doctors of Nigeria.* **16**(4):364-7, 2007 Oct-Dec.

Uludag S, Salihoglu SF, Madazli R, Cepni I. A comparison of oral and vaginal misoprostol for induction of labor. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2005;**122**:57-60.

Urban R, Lemancewicz A, Urban J, Skotnicki MZ, Kretowska M. Misoprostol and dinoprostone therapy for labor induction: a Doppler comparison of uterine and fetal hemodynamic effects. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2003;**106**:20-4.

van Gemund N, Scherjon S, LeCessie S, van Leeuwen JH, van Roosmalen J, Kanhai HH. A randomised trial comparing low dose vaginal misoprostol and dinoprostone for labour induction. *BJOG.* 2004;**111**:42-9.

Varaklis K, Gumina R, Stubblefield PG. Randomized controlled trial of vaginal misoprostol and intracervical prostaglandin E2 gel for induction of labor at term. *Obstet.Gynecol.* 1995;**86**:541-4.

Vengalil SR, Guinn DA, Olabi NF, Burd LI, Owen J. A randomized trial of misoprostol and extra-amniotic saline infusion for cervical ripening and labor induction. *Obstet.Gynecol.* 1998;**91**:774-9.

Vengalil SR, Guinn DA, Olabi NF, Burd LI, Owen J. Randomized trial of misoprostol and extraamniotic saline infusion for cervical ripening and labor induction. *Obstetrics and Gynecology* 1998;**91**:774-9.

Wang H, Li L, Pu L. [The effect of 25 micrograms misoprostol on induction of labor in late pregnancy]. *Zhonghua Fu Chan Ke.Za Zhi.* 1998;**33**:469-71.

Wang L, Shi C, Yang G. [Comparison of misoprostol and ricinus oil meal for cervical ripening and labor induction]. *Zhonghua Fu Chan Ke.Za Zhi*. 1997;**32**:666-8.

Wang Z, Li W, Ouyang W. [Safety and efficacy of intravaginal misoprostol for cervical ripening in the third trimester of pregnancy]. *Zhonghua Fu Chan Ke.Za Zhi*. 1997;**32**:326-8.

Wang Z, Li W, Ouyang W, Ding Y, Wang F, Xu L *et al*. Cervical ripening in the third trimester of pregnancy with intravaginal misoprostol: a double-blind, randomized, placebo-controlled study. *J.Tongji Med.Univ* 1998;**18**:183-6.

Wilk M, Poreba R, Sipinski A, Jureczko T. [The use of Misoprostol preparation (Cytotec) in induction of labor at prolonged pregnancy]. *Ginekol.Pol*. 2000;**71**:322-6.

Wilk M, Jureczko T, Poreba R, Sipinski A. [Misoprostol and oxytocin in induction of labour in women with prolonged pregnancy--safety and effectiveness comparison]. *Wiad.Lek*. 2001;**54**:662-7.

Windrim R, Bennett K, Mundle W, Young DC. Oral administration of misoprostol for labor induction: a randomized controlled trial. *Obstet.Gynecol*. 1997;**89**:392-7.

Wing DA, Rahall A, Jones MM, Goodwin TM, Paul RH. Misoprostol: an effective agent for cervical ripening and labor induction. *Am.J.Obstet.Gynecol*. 1995;**172**:1811-6.

Wing DA, Jones MM, Rahall A, Goodwin TM, Paul RH. A comparison of misoprostol and prostaglandin E2 gel for preinduction cervical ripening and labor induction. *Am.J.Obstet.Gynecol*. 1995;**172**:1804-10.

Wing DA, Rahall A, Jones MM, Goodwin TM, Paul RH. Misoprostol - An Effective Agent for Cervical Ripening and Labor Induction. *American Journal of Obstetrics and Gynecology* 1995;**172**:1811-6.

Wing DA, Paul RH. A comparison of differing dosing regimens of vaginally administered misoprostol for preinduction cervical ripening and labor induction. *Am.J.Obstet.Gynecol*. 1996;**175**:158-64.

Wing DA, Ortiz-Omphroy G, Paul RH. A comparison of intermittent vaginal administration of misoprostol with continuous dinoprostone for cervical ripening and labor induction. *Am.J.Obstet.Gynecol*. 1997;**177**:612-8.

Wing DA, Lovett K, Paul RH. Disruption of prior uterine incision following misoprostol for labor induction in women with previous cesarean delivery. *Obstet.Gynecol*. 1998;**91**:828-30.

Wing DA, Paul RH. Induction of labor with misoprostol for premature rupture of membranes beyond thirty-six weeks' gestation. *Am.J.Obstet.Gynecol*. 1998;**179**:94-9.

- Wing DA, Ham D, Paul RH. A comparison of orally administered misoprostol with vaginally administered misoprostol for cervical ripening and labor induction. *Am.J.Obstet.Gynecol.* 1999;**180**:1155-60.
- Wing DA, Fassett MJ, Mishell DR. Mifepristone for preinduction cervical ripening beyond 41 weeks' gestation: a randomized controlled trial. *Obstet.Gynecol.* 2000;**96**:543-8.
- Wing DA, Park MR, Paul RH. A randomized comparison of oral and intravaginal misoprostol for labor induction. *Obstet.Gynecol.* 2000;**95**:905-8.
- Wing DA, Fassett MJ, Guberman C, Tran S, Parrish A, Guinn D. A comparison of orally administered misoprostol to intravenous oxytocin for labor induction in women with favorable cervical examinations. *Am.J.Obstet.Gynecol.* 2004;**190**:1689-94.
- Wing DA, Guberman C, Fassett M. A randomized comparison of oral mifepristone to intravenous oxytocin for labor induction in women with prelabor rupture of membranes beyond 36 weeks' gestation. *American Journal of Obstetrics and Gynecology* 2005;**192**:445-51.
- Wolf SB, Sanchez-Ramos L, Kaunitz AM. Sublingual misoprostol for labor induction: a randomized clinical trial. *Obstet.Gynecol.* 2005;**105**:365-71.
- Yin CY, Zhou JZ, Wang BP, Lu XY. [Effect and risk analysis of misoprostol in stimulating cervical maturity for post-term pregnancy]. *Nan.Fang Yi.Ke.Da.Xue.Xue.Bao.* 2006;**26**:182-4, 188.
- Young D, Delaney T, Armson T, Fanning C. Lower dose vaginal and oral misoprostol in labor induction - RCT. *American Journal of Obstetrics and Gynecology* 2001;**185**:S203.
- Zeteroglu S, Sahin HG, Sahin HA. Induction of labor with misoprostol in grand multiparous patients. *Int.J.Gynaecol.Obstet.* 2004;**87**:155-6.
- Zeteroglu S, Engin-Ustun Y, Ustun Y, Guvercinci M, Sahin G, Kamaci M. A prospective randomized study comparing misoprostol and oxytocin for premature rupture of membranes at term. *J.Matern.Fetal Neonatal Med.* 2006;**19**:283-7.
- Zeteroglu S, Sahin HG, Sahin HA. Induction of labor in great grandmultipara with misoprostol. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2006;**126**:27-32.
- Zeteroglu S, Sahin GH, Sahin HA. Induction of labor with misoprostol in pregnancies with advanced maternal age. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2006;**129**:140-4.
- Tanir HM, Sener T, Yildiz C, Kaya M, Kurt I. A prospective randomized trial of labor induction with vaginal controlled-release dinoprostone inserts with or without oxytocin and misoprostol+oxytocin. *Clinical & Experimental Obstetrics & Gynecology.* **35**(1):65-8, 2008.

Abdul MA, Ibrahim UN, Yusuf MD, Musa H. Efficacy and safety of misoprostol in induction of labour in a Nigerian tertiary hospital. *West African Journal of Medicine*. 2007 Jul-Sep; **26**(3):213-6.

Observational Studies

[Anon]. Misoprostol for cervical ripening: Is it a safe, cost-effective alternative to dinoprostone? *Formulary* 1999; **34**:77-8.

Aboufalah A, Chraibi T, el Mouatacim K, Samouh N, Himmi A. Induction of labour with intravaginal misoprostol after prior cesarean delivery. *Afr.J.Reprod.Health* 2001; **5**:139-42.

Akhan SE, Iyibozkurt AC, Turfanda A. Unscarred uterine rupture after induction of labor with misoprostol: a case report. *Clin.Exp.Obstet.Gynecol*. 2001; **28**:118-20.

Andresen DM, Jensen JS, Uldbjerg N. [Misoprostol--a safe preparation for induction of labor?]. *Ugeskr.Laeger* 2006; **168**:3711-4.

Ara J, Noorani M. Induction of labour with oral misoprostol for prelabour rupture of membranes at term. *J.Pak.Med.Assoc*. 2005; **55**:180-3.

Aslan H, Unlu E, Agar M, Ceylan Y. Uterine rupture associated with misoprostol labor induction in women with previous cesarean delivery. *Eur.J.Obstet.Gynecol.Reprod.Biol*. 2004; **113**:45-8.

Bennett BB. Uterine rupture during induction of labor at term with intravaginal misoprostol. *Obstet.Gynecol*. 1997; **89**:832-3.

Berkley E, Meng C, Rayburn WF. Success rates with low dose misoprostol before induction of labor for nulliparas with severe preeclampsia at various gestational ages. *Journal of Maternal-Fetal & Neonatal Medicine*. 20(11):825-31, 2007 Nov.

Bique C, Bugalho A, Bergstrom S. Labor induction by vaginal misoprostol in grand multiparous women. *Acta Obstet.Gynecol.Scand*. 1999; **78**:198-201.

Blanchette HA, Nayak S, Erasmus S. Comparison of the safety and efficacy of intravaginal misoprostol (prostaglandin E1) with those of dinoprostone (prostaglandin

- E2) for cervical ripening and induction of labor in a community hospital. *Am.J.Obstet.Gynecol.* 1999;**180**:1551-9.
- Bugalho A, Bique C, Machungo F, Bergstrom S. A comparative study of vaginal misoprostol and intravenous oxytocin for induction of labour. *Gynecol.Obstet.Invest* 1995;**39**:252-6.
- Bugalho A, Bique C, Machungo F, Faundes A. Low-dose vaginal misoprostol for induction of labor with a live fetus. *Int.J.Gynaecol.Obstet.* 1995;**49**:149-55.
- Buhimschi CS, Buhimschi IA, Patel S, Malinow AM, Weiner CP. Rupture of the uterine scar during term labour: contractility or biochemistry? *BJOG.* 2005;**112**:38-42.
- Bush MC, Csaba A, Eddleman KA, Saphier CJ. Is misoprostol safe for labor induction in twin gestations? *J.Matern.Fetal Neonatal Med.* 2006;**19**:35-8.
- Caliskan E, Dilbaz S, Gelisen O, Dilbaz B, Ozturk N, Haberal A. Unsuccessful labour induction in women with unfavourable cervical scores: predictors and management. *Aust.N.Z.J.Obstet.Gynaecol.* 2004;**44**:562-7.
- Castaneda CS, Izquierdo Puente JC, Leon Ochoa RA, Plasse TF, Powers BL, Rayburn WF. Misoprostol dose selection in a controlled-release vaginal insert for induction of labor in nulliparous women. *Am.J.Obstet.Gynecol.* 2005;**193**:1071-5.
- Chandra S, Allen V, Lee W, Fanning C, Young D. The effect of vaginal pH on labor induction with vaginal misoprostol. *J.Matern.Fetal Neonatal Med.* 2005;**17**:387-91.
- Chang DW, Velazquez MD, Colyer M, Klaus P, Mallipeddi SK, Rayburn WF. Vaginal misoprostol for cervical ripening at term: comparison of outpatient vs. inpatient administration. *J.Reprod.Med.* 2005;**50**:735-9.
- Chen WH, Lai HC, Tang YH, Liu HS. Fetal Doppler hemodynamic changes in spontaneous versus prostaglandin E-1-induced active labor. *Acta Obstetricia et Gynecologica Scandinavica* 1999;**78**:599-604.
- Cheng SY, Chen TC. Pilot study of labor induction with titrated oral misoprostol. *Taiwan.J.Obstet.Gynecol.* 2006;**45**:225-9.
- Choy-Hee L, Raynor BD. Misoprostol induction of labor among women with a history of cesarean delivery. *Am.J.Obstet.Gynecol.* 2001;**184**:1115-7.
- Chraibi T, Aboufalah A, Bisbis W, Noun M, Matar N, Samouh N *et al.* [Induction of labor by misoprostol, an analog of PGE1. A prospective study of 200 cases]. *Sante* 1999;**9**:345-9.
- Clark S, Blum J, Blanchard K, Galvao L, Fletcher H, Winikoff B. Misoprostol use in obstetrics and gynecology in Brazil, Jamaica, and the United States. *Int.J.Gynaecol.Obstet.* 2002;**76**:65-74.

Crane JM, Young DC, Butt KD, Bennett KA, Hutchens D. Excessive uterine activity accompanying induced labor. *Obstet.Gynecol.* 2001;**97**:926-31.

Crane JM, Delaney T, Butt KD, Bennett KA, Hutchens D, Young DC. Predictors of successful labor induction with oral or vaginal misoprostol. *J.Matern.Fetal Neonatal Med.* 2004;**15**:319-23.

Dadi H, Patwardhan SC, Ikomi A, Ojutiku D. Self-induction of labour using misoprostol: an unusual case of uterine hyperstimulation. *BJOG.* 2004;**111**:1018-9.

Daisley H. Case reports 1. Maternal mortality following the use of misoprostol. *Medicine Science and the Law* 2000;**40**:78-82.

Daisley H, Jr. Maternal mortality following the use of misoprostol. *Med.Sci.Law* 2000;**40**:78-82.

Dede FS, Haberal A, Dede H, Sivaslioglu A, Arslanpence I. Misoprostol for cervical ripening and labor induction in pregnancies with oligohydramnios. *Gynecol.Obstet.Invest* 2004;**57**:139-43.

Del Valle GO, Sanchez-Ramos L, Jordan CW, Gaudier FL, Delke I. Use of misoprostol (prostaglandin E1 methyl analogue) to expedite delivery in severe preeclampsia remote from term. *J.Matern.Fetal Med.* 1996;**5**:39-40.

Delaney T, Crane JM, Hutchens D, Fanning CA, Young DC. Oral misoprostol labor induction in patients with a favorable cervix. *American Journal of Obstetrics and Gynecology* 2001;**185**:S202.

Diab AE. Uterine ruptures in Yemen. *Saudi.Med.J.* 2005;**26**:264-9.

Ding DC, Su HY, Chu TW. Intravaginal and intracervical misoprostol for cervical ripening of labor in primiparas. *Int.J.Gynaecol.Obstet.* 2003;**82**:209-11.

Ding DC, Hsu S, Su HY. Low dose intravaginal misoprostol for induction of labor at term. *Int.J.Gynaecol.Obstet.* 2005;**90**:72-3.

Escumalha M, Gouveia C, Cunha M, Vale F, Machado MC. Neonatal morbidity and outcome of live born premature babies after attempted illegal abortion with misoprostol. *Pediatr.Nurs.* 2005;**31**:228-31.

Ezechi OC, Kalu BK, Njokanma FO, Nwokoro CA, Okeke GC. Vaginal misoprostol induction of labour: a Nigerian hospital experience. *J.Obstet.Gynaecol.* 2004;**24**:239-42.

Ezegwui HU. Uterine rupture in a primigravida when misoprostol was used for induction of labour and subsequent successful pregnancy outcome. *J.Obstet.Gynaecol.* 2006;**26**:160-1.

- Feitosa FE, de Amorim MM, Alencar Junior CA, Coutinho IC, Sampaio ZS. [New formulation of sublingual misoprostol (25 mcg) for induction of labor]. *Rev.Assoc.Med.Bras.* 2006;**52**:251-5.
- Fittkow CT, Maul H, Olson G, Martin E, MacKay LB, Saade GR *et al.* Light-induced fluorescence of the human cervix decreases after prostaglandin application for induction of labor at term. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2005;**123**:62-6.
- Fletcher H.,McCaw-Binns A. Rupture of the uterus with misoprostol (prostaglandin E1) used for induction of labour. *J.Obstet.Gynaecol.* 1998;**18**:184-5.
- Fletcher H.,Hutchinson S. A retrospective review of pregnancy outcome after misoprostol (prostaglandin E1) induction of labour. *West Indian Med.J.* 2001;**50**:47-9.
- Fontenot MT, Lewis DF, Barton CB, Jones EM, Moore JA, Evans AT. Abruptio placentae associated with misoprostol use in women with preeclampsia. *J.Reprod.Med.* 2005;**50**:653-8.
- Gherman RB, McBrayer S, Browning J. Uterine rupture associated with vaginal birth after cesarean section: a complication of intravaginal misoprostol? *Gynecol.Obstet.Invest* 2000;**50**:212-3.
- Gilson G, Burt A, Holck P, Murray K, Murphy N. Outpatient preinduction cervical ripening with low-dose intravaginal misoprostol. *American Journal of Obstetrics and Gynecology* 2003;**189**:S200.
- Godinjak Z, Imsirija L, Imsirija N. [Vaginal application of misoprostol for labor induction after 36 weeks of pregnancy]. [Bosnian] *Medicinski Arhiv.* **61**(4):218-20, 2007.
- Goffinet F, Humbert R, Clerson P, Philippe HJ, Breart G, Cabrol D. [National survey on the use of induced labor by obstetricians. Study Group on Induced Labor]. *J.Gynecol.Obstet.Biol.Reprod.(Paris)* 1999;**28**:319-29.
- Gunalp S.,Bildirici I. The effect of vaginal pH on the efficacy of vaginal misoprostol for induction of labor. *Acta Obstet.Gynecol.Scand.* 2000;**79**:283-5.
- Hascalik S, Celik O, Dogru MI, Kocagun DA, Inan E, Yurekli M. Influence of misoprostol (PGE1) on amniotic fluid and maternal serum adrenomedullin levels. *Acta Obstet.Gynecol.Scand.* 2005;**84**:833-6.
- Hendrix NW, Chauhan SP, Morrison JC, Magann EF, Martin JN, Devoe LD. Bishop score: A poor diagnostic test to predict failed induction versus vaginal delivery. *Southern Medical Journal* 1998;**91**:248-52.
- Hill DA, Chez RA, Quinlan J, Fuentes A, LaCombe J. Uterine rupture and dehiscence associated with intravaginal misoprostol cervical ripening. *J.Reprod.Med.* 2000;**45**:823-6.

- Hoffman MK, Sciscione A, Srinivasana M, Shackelford DP, Ekbladh L. Uterine rupture in patients with a prior Cesarean delivery: The impact of cervical ripening. *American Journal of Perinatology* 2004;**21**:217-22.
- Hoffman MK, Vahratian A, Sciscione AC, Troendle JF, Zhang J. Comparison of Labor progression between induced and noninduced multiparous women. *Obstetrics and Gynecology* 2006;**107**:1029-34.
- Hofmeyr GJ, Matonhodze BB, Alfirevic Z, Campbell E, de Jager M, Nikodem C. Titrated oral misoprostol solution--a new method of labour induction. *S.Afr.Med.J.* 2001;**91**:775-6.
- Hsieh YY, Tsai HD, Chang CC, Yeh LS, Yang TC, Hsu TY. Precipitate delivery and postpartum hemorrhage after term induction with 200 micrograms misoprostol. *Zhonghua Yi.Xue.Za Zhi.(Taipei)* 2000;**63**:58-61.
- Jouatte F, Subtil D, Marquis P, Plennevaux JL, Puech F. [Medical indications of labor induction: a comparison between intravaginal misoprostol and intravenous prostaglandin E2]. *J.Gynecol.Obstet.Biol.Reprod.(Paris)* 2000;**29**:763-71.
- Katz VL, Farmer RM, Dean CA, Carpenter ME. Use of misoprostol for cervical ripening. *South.Med.J.* 2000;**93**:881-4.
- Khosla AH, Sirohiwal D, Sangwan K. A still birth and uterine rupture during induction of labour with oral misoprostol. *Aust.N.Z.J.Obstet.Gynaecol.* 2002;**42**:412.
- Kimya Y, Ozerkan K, Karabay A. Dehydroepiandrosterone sulfate levels and success of labor induction. *Int.J.Gynaecol.Obstet.* 2004;**86**:33-4.
- Lapaire O, Zanetti-Dallenbach R, Weber P, Hosli I, Holzgreve W, Surbek D. Labor induction in preeclampsia: is misoprostol more effective than dinoprostone? *J.Perinat.Med.* 2007;**35**:195-9.
- Levey KA, MacKenzie AP, Stephenson C, Bercik R, Kuczynski E, Funai EF. Increased rates of chorioamnionitis with extra-amniotic saline infusion method of labor induction. *Obstet.Gynecol.* 2004;**103**:724-8.
- Levey KA, Arslan AA, Funai EF. Extra-amniotic saline infusion increases cesarean risk versus other induction methods and spontaneous labor. *American Journal of Perinatology* 2006;**23**:435-8.
- Li L, Gao W, Chen S. [Labour induction in women at term with mifepristone and misoprostol]. *Zhonghua Fu Chan Ke.Za Zhi.* 1996;**31**:681-4.
- Lieberman E. Risk factors for uterine rupture during a trial of labor after cesarean. *Clinical Obstetrics and Gynecology* 2001;**44**:609-21.

- Lin C.,Raynor BD. Risk of uterine rupture in labor induction of patients with prior cesarean section: an inner city hospital experience. *Am.J.Obstet.Gynecol.* 2004;**190**:1476-8.
- Liu HS, Chu TY, Chang YK, Yu MH, Chen WH. Intracervical misoprostol as an effective method of labor induction at term. *Int.J.Gynaecol.Obstet.* 1999;**64**:49-53.
- Loto OM, Fadahunsi AA, Kolade CO. Safety and efficacy of misoprostol for induction of labour in a semi-urban hospital setting. *J.Obstet.Gynaecol.* 2004;**24**:638-40.
- Lydon-Rochelle M, Holt VL, Easterling TR, Martin DP. Risk of uterine rupture during labor among women with a prior cesarean delivery. *New England Journal of Medicine* 2001;**345**:3-8.
- Majoko F, Magwali T, Zwizwai M. Uterine rupture associated with use of misoprostol for induction of labor. *Int.J.Gynaecol.Obstet.* 2002;**76**:77-8.
- Margulies M, Catuzzi P, Voto LS, Imaz FU. Induction of Labor with A Pge1 Analog - An Open, Noncomparative, Dose Finding Study. *Prensa Medica Argentina* 1991;**78**:9-13.
- Mathews JE, Mathai M, George A. Uterine rupture in a multiparous woman during labor induction with oral misoprostol. *Int.J.Gynaecol.Obstet.* 2000;**68**:43-4.
- Mazzone ME.,Woolever J. Uterine rupture in a patient with an unscarred uterus: a case study. *WMJ.* 2006;**105**:64-6.
- Mbele AM, Makin JD, Pattinson RC. Can the outcome of induction of labour with oral misoprostol be predicted. *S.Afr.Med.J.* 2007;**97**:289-92.
- McGill J.,Shetty A. Mifepristone and misoprostol in the induction of labor at term. *Int.J.Gynaecol.Obstet.* 2007;**96**:80-4.
- McKenna DS.,Duke JM. Effectiveness and infectious morbidity of outpatient cervical ripening with a Foley catheter. *Journal of Reproductive Medicine* 2004;**49**:28-32.
- Merrell DA.,Koch MA. Induction of labour with intravaginal misoprostol in the second and third trimesters of pregnancy. *S.Afr.Med.J.* 1995;**85**:1088-90.
- Mousa HA.,Mahmood TA. Do practice guidelines guide practice? A prospective audit of induction of labor - three years experience. *Acta Obstetricia et Gynecologica Scandinavica* 2000;**79**:1086-92.
- Mutlu MM, Caliskan E, Haberal A. Prediction of adverse outcome associated with vaginal misoprostol for labor induction. *Eur.J.Obstet.Gynecol.Reprod.Biol.* 2003;**110**:143-8.

- Oyelese Y, Landy HJ, Collea JV. Cervical laceration associated with misoprostol induction. *Int.J.Gynaecol.Obstet.* 2001;**73**:161-2.
- Ozan H, Uncu G, Yildirim V, Omak M, Kara HF, Tufekci M. Misoprostol in labor induction. *J.Obstet.Gynaecol.Res.* 2001;**27**:17-20.
- Ozden S, Delikara MN, Avci A, Ficicioglu C. Intravaginal misoprostol vs. expectant management in premature rupture of membranes with low Bishop scores at term. *Int.J.Gynaecol.Obstet.* 2002;**77**:109-15.
- Pajak J, Tomialowicz M, Florjanski J, Heimrath J, Myszczyzyn G, Zalewski J *et al.* [Comparison of vaginal misoprostol and oxytocin for labor induction in post-term pregnancy]. *Ginekol.Pol.* 2001;**72**:1300-4.
- Parer J, Nageotte M, Rowles R, Schwartz M, Barford D, Garite T *et al.* Factors affecting the likelihood of successful induction after intravaginal misoprostol application for cervical ripening and labor induction - Discussion. *American Journal of Obstetrics and Gynecology* 2002;**186**:1240-3.
- Phillip H, Fletcher H, Reid M. The impact of induced labour on postpartum blood loss. *J.Obstet.Gynaecol.* 2004;**24**:12-5.
- Plaut MM, Schwartz ML, Lubarsky SL. Uterine rupture associated with the use of misoprostol in the gravid patient with a previous cesarean section. *Am.J.Obstet.Gynecol.* 1999;**180**:1535-42.
- Politz LB, Chez RA, Parsons MT, Huffman K. The effect of a pill inserter on vaginal misoprostol dosing. *J.Matern.Fetal Med.* 2001;**10**:332-4.
- Raio L, Ghezzi F, Di Naro E, Bruhwiler H, Luscher KP. Shorter delivery time after labor induction with misoprostol. *Zeitschrift fur Geburtshilfe und Neonatologie* 2001;**205**:147-51.
- Ramsey PS, Ogburn PL, Jr., Harris DY, Heise RH, Ramin KD. Effect of vaginal pH on efficacy of misoprostol for cervical ripening and labor induction. *Am.J.Obstet.Gynecol.* 2000;**182**:1616-9.
- Rayburn WF, Sorkary N, Clokey DE, Moore LE, Curet LB. Consequences of routine delivery at 38 weeks for A-2 gestational diabetes. *J.Matern.Fetal Neonatal Med.* 2005;**18**:333-7.
- Rayburn WF, Powers BL, Plasse TF, Carr D, Di Spirito M. Pharmacokinetics of a controlled-release misoprostol vaginal insert at term. *J.Soc.Gynecol.Investig.* 2006;**13**:112-7.
- Rodrigues R, Nunes F, Tiago D, Avillez T, Vieira A, Meirinho M. Induction of labor with intravaginal administration of misoprostol. *Int.J.Gynaecol.Obstet.* 1998;**60**:233-7.

Rozenberg P, Chevret S, Chastang C, Ville Y. Comparison of digital and ultrasonographic examination of the cervix in predicting time interval from induction to delivery in women with a low Bishop score. *BJOG*. 2005;**112**:192-6.

Sachs BP. A 38-year-old woman with fetal loss and hysterectomy. *JAMA* 2005;**294**:833-40.

Sahin HG, Sahin HA, Surucu R, Guvercinci M. A study of intravaginal misoprostol for induction of labor in toxemia of pregnancy. *Int.J.Gynaecol.Obstet.* 2001;**75**:3-9.

Samuels TA, Awonuga A. Second-trimester rudimentary uterine horn pregnancy: rupture after labor induction with misoprostol. *Obstet.Gynecol.* 2005;**106**:1160-2.

Sanchez-Ramos L, Kaunitz AM. Perinatal morbidity associated with misoprostol-induced labor. *Am.J.Obstet.Gynecol.* 2000;**182**:749-50.

Santana-Casteneda C, Izquierdo-Puente JC, Leon-Ochoa RA, Plasse T, Rayburn W. Misoprostol controlled-release vaginal insert for cervical ripening and induction of labor in nulliparous women. *American Journal of Obstetrics and Gynecology* 2004;**191**:S187.

Schroder AK, Tauchert S, Diedrich K. [Induction of labour at term with misoprostol: an effective, safe and inexpensive alternative]. *Zentralbl.Gynakol.* 2004;**126**:154-8.

Sciscione A, Larkin M, O'Shea A, Pollock M, Hoffman M, Colmorgen G. Preinduction cervical ripening with the Foley catheter and the risk of subsequent preterm birth. *American Journal of Obstetrics and Gynecology* 2004;**190**:751-4.

Sciscione AC, Nguyen L, Manley JS, Shlossman PA, Colmorgen GH. Uterine rupture during preinduction cervical ripening with misoprostol in a patient with a previous Caesarean delivery. *Aust.N.Z.J.Obstet.Gynaecol.* 1998;**38**:96-7.

Shetty A, Burt R, Rice P, Templeton A. Women's perceptions, expectations and satisfaction with induced labour - A questionnaire-based study. *European Journal of Obstetrics Gynecology and Reproductive Biology* 2005;**123**:56-61.

Simoes T, Condeco P, Dias E, Ventura P, Matos C, Blickstein I. Induction of labor with oral misoprostol in nulliparous mothers of twins. *J.Perinat.Med.* 2006;**34**:111-4.

Singh BM, Kriplani A, Bhatla N. Uterocervical laceration during induction of labour with intravaginal misoprostol in a woman with idiopathic thrombocytopenic purpura. *J.Obstet.Gynaecol.* 2005;**25**:75-6.

Szczesny W, Kjollesdal M, Karlsson B, Nielsen S. Bishop score and the outcome of labor induction with misoprostol. *Acta Obstet.Gynecol.Scand.* 2006;**85**:579-82.

Thomas A, Jophy R, Maskhar A, Thomas RK. Uterine rupture in a primigravida with misoprostol used for induction of labour. *BJOG*. 2003;**110**:217-8.

Vahratian A, Zhang J, Troendle JF, Sciscione AC, Hoffman MK. Labor progression and risk of cesarean delivery in electively induced nulliparas. *Obstetrics and Gynecology* 2005;**105**:698-704.

Wagner M. Misoprostol (Cytotec) for labor induction: a cautionary tale. *Midwifery Today Int.Midwife*. 1999;31-3.

Wagner M. Midwives and cytotec: a true story. *Midwifery Today Int.Midwife*. 2001;44-5.

Ware V, Raynor BD. Transvaginal ultrasonographic cervical measurement as a predictor of successful labor induction. *Am.J.Obstet.Gynecol*. 2000;**182**:1030-2.

Wing DA, Tran S, Paul RH. Factors affecting the likelihood of successful induction after intravaginal misoprostol application for cervical ripening and labor induction. *Am.J.Obstet.Gynecol*. 2002;**186**:1237-40.

Yogev Y, Ben Haroush A, Lahav E, Horowitz E, Hod M, Kaplan B. Induction of labor with prostaglandin E-2 in women with previous cesarean section and unfavorable cervix. *European Journal of Obstetrics Gynecology and Reproductive Biology* 2004;**116**:173-6.

Yolande H, Namory K, Delphine F, Mamadou HD, Mamadou DB, Daniel T *et al*. Misoprostol use for labor induction in developing countries: a prospective study in Guinea. *Eur.J.Obstet.Gynecol.Reprod.Biol*. 2005;**122**:40-4.

Ekele BA, Nnadi DC, Gana MA, Shehu CE, Ahmed Y, Nwobodo EI. Misoprostol use for cervical ripening and induction of labour in a Nigerian teaching hospital. *Nigerian Journal of Clinical Practice*. **10**(3):234-7, 2007 Sep.

Cost-effectiveness

Ramsey PS, Harris DY, Ogburn PL, Heise RH, Magtibay PM, Ramin KD. Comparative cost analysis of prostaglandin analogues dinoprostone and misoprostol as labor preinduction agents. *Prim.Care Update.Ob Gyns*. 1998;**5**:182.

Ramsey PS, Harris DY, Ogburn PL, Jr., Heise RH, Magtibay PM, Ramin KD. Comparative efficacy and cost of the prostaglandin analogs dinoprostone and misoprostol as labor preinduction agents. *Am.J.Obstet.Gynecol*. 2003;**188**:560-5.

Taylor SJ, Armour CL. Cost implications in the management of induction of labour. *Pharmacoeconomics* 1997;**12**:547-54.

Pharmacology

Alan M, Tasal I. Efficacy of prostaglandin F2alpha and misoprostol in the induction of parturition in goats. *Vet.Rec.* 2002;**150**:788-9.

Goldberg AB, Greenberg MB, Darney PD. Misoprostol and pregnancy. *N.Engl.J.Med.* 2001;**344**:38-47.

Hemmerling A. The safety of misoprostol. *International Journal of Gynecology & Obstetrics* 2006;**94**:S149-S150.

Hofmeyr GJ, Matonhodze BB, Alfirevic Z, Campbell E, de Jager M, Nikodem C. Titrated oral misoprostol solution--a new method of labour induction. *S.Afr.Med.J.* 2001;**91**:775-6.

Lebel W, Riccardi K, Grasser WA, Terry K, Thompson D, Paralkar VM. Prostaglandin E2 receptor subtype EP-2 is not involved in the induction of non-pregnant guinea pig uterine contractions associated with terminal pregnancy. *Prostaglandins Leukot.Essent.Fatty Acids* 2004;**71**:399-404.

Lyons C, Beharry K, Akmal Y, Attenello F, Nageotte MP. In vitro response of prostaglandin E2 receptor (EP3) in the term pregnant rat uterus and cervix to misoprostol. *Prostaglandins Other Lipid Mediat.* 2003;**70**:317-29.

Matonhodze BB, Katsoulis LC, Hofmeyr GJ. Labor induction and meconium: in vitro effects of oxytocin, dinoprostone and misoprostol on rat ileum relative to myometrium. *J.Perinat.Med.* 2002;**30**:405-10.

Rayburn WF, Powers BL, Plasse TF, Carr D, Di Spirito M. Pharmacokinetics of a controlled-release misoprostol vaginal insert at term. *J.Soc.Gynecol.Investig.* 2006;**13**:112-7.

Schaff EA, DiCenzo R, Fielding SL. Comparison of misoprostol plasma concentrations following buccal and sublingual administration. *Contraception* 2005;**71**:22-5.

Sitruk-Ware R. Mifepristone and misoprostol sequential regimen side effects, complications and safety. *Contraception* 2006;**74**:48-55.

Smith GCS, Baguma-Nibasheka M, Wu WX, Nathanielsz PW. Regional variations in contractile responses to prostaglandins and prostanoid receptor messenger ribonucleic acid in pregnant baboon uterus. *American Journal of Obstetrics and Gynecology* 1998;**179**:1545-52.

Wagner M. Adverse events following misoprostol induction of labor. *Midwifery Today Int.Midwife*. 2004;9-12.

Zhang WS, Chen LX, Wu ZD, Liu JX. [Effects of oxytocin, misoprostol and nimodipine on expression of L-type voltage dependent calcium channel mRNA of the uterine myometrium and left ventricular myocardial cells in late pregnant rats]. *Zhonghua Fu Chan Ke.Za Zhi*. 2003;**38**:1-4.

Zust S, Hosli I, Surbek D, Holzgreve W. Side effects of Misoprostol therapy during pregnancy. *Zeitschrift fur Geburtshilfe und Neonatologie* 2001;**205**:43-8.

Zust S, Hosli I, Surbek D, Holzgreve W. [Adverse effects of misoprostol in pregnancy]. *Z.Geburtshilfe Neonatol*. 2001;**205**:43-8.

Politics/ Licensing/ Ethics

ACOG Committee Opinion. Number 283, May 2003. New U.S. Food and Drug Administration labeling on Cytotec (misoprostol) use and pregnancy. *Obstet.Gynecol*. 2003;**101**:1049-50.

[Anon]. Urgent need for standardised misoprostol regimens. *Reproductive Health Matters* 2002;**10**:202.

[Anon]. USFDA, pharmacia new ruling on misoprostol. *Reproductive Health Matters* 2002;**10**:192-3.

[Anon]. New US Food and Drug Administration labeling on Cytotec (Misoprostol) use and pregnancy. *International Journal of Gynecology & Obstetrics* 2003;**82**:137-8.

Chong YS, Su LL, Arulkumaran S. Misoprostol: a quarter century of use, abuse, and creative misuse. *Obstet.Gynecol.Surv*. 2004;**59**:128-40.

Gebhardt DO. Misoprostol in a topsyturvy world. *J.Med.Ethics* 2001;**27**:205.

Goer H. Cytotec (misoprostol) - Reply. *Birth-Issues in Perinatal Care* 2003;**30**:285.

Howarth GR, Pistorius L, Mantel G, Funk M, Pattinson RC. Induction of labour at term--misoprostol, efficacy, economics and ethics. *S.Afr.Med.J*. 1996;**86**:1174, 1176.

Mackenzie WE. Misoprostol and the politics of fear. *Lancet* 2001;**357**:1296.

Wagner M. Misoprostol and the politics of convenience. *Lancet* 2001;**357**:2142.

Wagner M. Off-label use of misoprostol in obstetrics: a cautionary tale. *BJOG*. 2005;**112**:266-8.

Weeks AD, Fiala C, Safar P. Misoprostol and the debate over off-label drug use. *BJOG*. 2005;**112**:269-72.

Editorials/ Letters/ Opinions

Butt K, Crane J, Young D, Bennett K, Hutchens D. Comparison of oral misoprostol and oxytocin for labor induction - Reply. *Obstetrics and Gynecology* 2000;**95**:786.

Carlan SJ. Safety and efficacy of misoprostol orally and vaginally: A randomized trial - Reply. *Obstetrics and Gynecology* 2001;**98**:876.

Crane JM, Young D, Butt K, Delaney M, Hutchens D. Safety and efficacy of misoprostol orally and vaginally: a randomized trial. *Obstet.Gynecol.* 2001;**98**:875-6.

Cullen M. Use of Cytotec (misoprostol) for labor induction. *J.Obstet.Gynecol.Neonatal Nurs.* 2001;**30**:11.

Day L, Tailor A, Howard R, Burrell S. Misoprostol for induction of labour at term. *BJOG*. 2000;**107**:576-7.

Devitt NF. Cytotec (misoprostol). *Birth* 2003;**30**:285.

Eaker JR, Wade RV, Hopwood HG, Parker L, Brown EH, Farah. Randomized trial of two doses of the prostaglandin E-1 analog misoprostol for labor induction - Discussion. *American Journal of Obstetrics and Gynecology* 1997;**177**:369-71.

Feingold M, Ghamande S. A randomized trial of misoprostol and oxytocin for induction of labor: safety and efficacy. *Obstet.Gynecol.* 1997;**90**:155.

Ferroni KM. Induction of labor: a clinician's viewpoint. *Ann.Pharmacother.* 1997;**31**:1401-2.

Fletcher H. Intravaginal Misoprostol Versus Dinoprostone As Cervical Ripening and Labor-Inducing Agents - Reply. *Obstetrics and Gynecology* 1994;**83**:800-1.

Gaskin IM. The problem with Cytotec is that it has not been approved for obstetrical use. *Birth Gaz.* 1999;**15**:3.

Gebhardt DO. Misoprostol in a topsyturvy world. *J.Med.Ethics* 2001;**27**:205.

Goer H. Cytotec (misoprostol) - Reply. *Birth-Issues in Perinatal Care* 2003;**30**:285.

Gonzalez Pena JC. Misoprostol for induction of labor. *Am.J.Obstet.Gynecol.* 1996;**174**:797.

- Greene MF. Vaginal delivery after cesarean section - Is the risk acceptable? *New England Journal of Medicine* 2001;**345**:54-5.
- Hale RW, Zinberg S. Use of misoprostol in pregnancy. *N.Engl.J.Med.* 2001;**344**:59-60.
- Hidar S, Khairi H. Outpatient misoprostol cervical ripening without subsequent induction of labor to prevent post-term pregnancy. *Acta Obstet.Gynecol.Scand.* 2005;**84**:1129.
- Hofmeyr GJ. Misoprostol in obstetrics and gynaecology--unregistered, dangerous and essential. *S.Afr.Med.J.* 1998;**88**:535-6.
- Hofmeyr GJ, Gulmezoglu AM, Alfirevic Z. Misoprostol for induction of labour at term - Reply. *British Journal of Obstetrics and Gynaecology* 2000;**107**:576-7.
- Irwin HW, Paper JT, Goodwin TM, Schroeder PW, Macer JA, Lee GF *et al.* Uterine rupture associated with the use of misoprostol in the gravid patient with a previous cesarean section - Discussion. *American Journal of Obstetrics and Gynecology* 1999;**180**:1540-2.
- Jackson N, Paterson-Brown S. Labour characteristics and uterine activity: misoprostol compared with oxytocin in women at term with prelabour rupture of the membranes. *BJOG.* 2000;**107**:1181-2.
- Kramer RL, Gilson GJ, Qualls CR. A randomized trial of misoprostol and oxytocin for induction of labor: Safety and efficiency - Reply. *Obstetrics and Gynecology* 1997;**90**:155.
- Laughead M. A comparison of orally administered misoprostol to intravenous oxytocin for labor induction in women with favorable cervical examinations - Discussion. *American Journal of Obstetrics and Gynecology* 2004;**190**:1694-5.
- Ledger WJ. Labor induction with intravaginal misoprostol in term premature rupture of membranes: a randomized study. *Obstet.Gynecol.* 1997;**90**:644.
- Lydon-Rochelle M, Holt VL, Easterling TR. Uterine rupture among women with a prior cesarean delivery - Reply. *New England Journal of Medicine* 2002;**346**:136.
- Mahajan NN. Induction of labor with oral misoprostol for premature rupture of membranes at term in women with unfavorable cervix. *J.Perinat.Med.* 2007.
- Majoko F. Misoprostol in obstetrics. *BJOG.* 2005;**112**:1666.
- McLean MT. Misoprostol. *Midwifery Today Int.Midwife.* 1999;5.
- Ngai SW, Chan YM, Lam H, Lao T. Labour characteristics and uterine activity: misoprostol compared with oxytocin in women at term with prelabour rupture of the

membranes - Authors' reply. *British Journal of Obstetrics and Gynaecology* 2000;**107**:1181-2.

O'Sullivan MJ, Frank F. Misoprostol versus low-dose oxytocin for cervical ripening: A prospective, randomized, double-masked trial - Discussion. *American Journal of Obstetrics and Gynecology* 2002;**187**:279-80.

Oboro V. Outpatient use of misoprostol in reducing gestation length - Reply. *Acta Obstetrica et Gynecologica Scandinavica* 2005;**84**:1130.

Prins R, Schwartz M, Girard J, Wing DA. A comparison of orally administered misoprostol to intravenous oxytocin for labor induction in women with favorable cervical examinations. *American Journal of Obstetrics and Gynecology* 2004;**190**:1695-6.

Sanchez-Ramos L, Kaunitz AM. Intravaginal misoprostol versus dinoprostone as cervical ripening and labor-inducing agents. *Obstet. Gynecol.* 1994;**83**:799-801.

Sanchez-Ramos L, Kaunitz AM. Oral administration of misoprostol for labor induction: a randomized controlled trial. *Obstet. Gynecol.* 1997;**90**:153-4.

Sanchez-Ramos L, Delke I. Induction of labor with misoprostol for premature rupture of membranes beyond thirty-six weeks' gestation. *Am. J. Obstet. Gynecol.* 1999;**180**:253-4.

Sanchez-Ramos L, Kaunitz AM. Uterine rupture associated with the use of prostaglandin E1 in patients with previous cesarean delivery. *Am. J. Obstet. Gynecol.* 2000;**182**:990-1.

Sanchez-Ramos L, Kaunitz AM. Outpatient cervical ripening with intravaginal misoprostol. *Obstet. Gynecol.* 2001;**97**:325-6.

Sanchez-Ramos L, Kaunitz AM. Labor induction with intravaginal misoprostol in term premature rupture of membranes: A randomized study - In reply. *Obstetrics and Gynecology* 1997;**90**:644.

Sciarra JJ. Announcing the winner of the IJGO Prize Paper Award for 2001. *Int. J. Gynaecol. Obstet.* 2002;**77**:105-7.

Stitely ML, Browning J, Gherman RB. Outpatient cervical ripening with intravaginal misoprostol - Reply. *Obstetrics and Gynecology* 2001;**97**:326.

Thulasimani M, Ramaswamy S. Comparison of oral misoprostol and oxytocin for labor induction. *Obstet. Gynecol.* 2000;**95**:786.

Wagner M. Misoprostol. *Midwifery Today Int. Midwife.* 1999;4-5.

Wing DA, Paul RH. Misoprostol for induction of labor - Reply. *American Journal of Obstetrics and Gynecology* 1996;**174**:797.

Wing DA, Paul RH. Induction of labor with misoprostol for premature rupture of membranes beyond thirty-six weeks' gestation - Reply. *American Journal of Obstetrics and Gynecology* 1999;**180**:253-4.

Young DC, Bennett K, Windrim R, Mundle W. Oral administration of misoprostol for labor induction: A randomized controlled trial - Reply. *Obstetrics and Gynecology* 1997;**90**:154.

Unspecified

Mother Jones magazine questions cytotec for labor induction. *AWHONN.Lifelines*. 2001;**5**:18-9.

Carvajal JA, Kushner L, Vera C, Oyarzun E. Combined misoprostol and oxytocin for cervical ripening and labor induction. *Journal of the Society for Gynecologic Investigation* 2004;**11**:163A.

Cunha M, Bugalho A, Bique C, Bergstrom S. Induction of labor by vaginal misoprostol in patients with previous cesarean delivery. *Acta Obstet.Gynecol.Scand.* 1999;**78**:653-4.

Duff P. Premature rupture of the membranes in term patients: Induction of labor versus expectant management. *Clinical Obstetrics and Gynecology* 1998;**41**:883-91.

Flamm BL. Vaginal birth after cesarean: Reducing medical and legal risks. *Clinical Obstetrics and Gynecology* 2001;**44**:622-9.

Flynn CA. Intravaginal misoprostol: a new option for labor induction? *J.Fam.Pract.* 1997;**44**:31.

How H, Ghulmiyyah L, Javaid S, Eruo F, Maxwell R, Recht K *et al.* Intravaginal misoprostol versus intravenous oxytocin for induction of labor in the preterm patient. *Journal of the Society for Gynecologic Investigation* 2006;**13**:121A-2A.

Howarth GR, Pistorius L, Mantel G, Funk M, Pattinson RC. Induction of labour at term--misoprostol, efficacy, economics and ethics. *S.Afr.Med.J.* 1996;**86**:1174, 1176.

Keating S, Kingdom J, Shannon P. Microscopic appearance of misoprostol following induction of labour. *Laboratory Investigation* 2004;**84**:271.

Koleva Z, Iankov M, Katsulov A, Savova S. [Induction of labor with misoprostol--vaginal administration]. *Akush.Ginekol.(Sofia)* 2005;**44**:10-2.

Mahlmeister L. Best practices in perinatal and neonatal nursing: cervical ripeners and the induction of labor. *J.Perinat.Neonatal Nurs.* 2005;**19**:97-9.

Mbele AM, Makin JD, Pattinson RC. Can the outcome of induction of labour with oral misoprostol be predicted. *South African Medical Journal. Suid-Afrikaanse Tydskrif Vir Geneeskunde.* **97**(4):289-92, 2007 Apr.

Mosler P, Necas M, Machac J. [Misoprostol and induction of labor--an effective method of treating prolonged pregnancy]. *Ceska.Gynekol.* 1998;**63**:325-6.

Powers BL, Rayburn WF, Wing D, Byrne J, Mercer B, Alfirevic Z. Safety profile of a controlled-release misoprostol vaginal insert for cervical ripening and labor induction. *Obstetrics and Gynecology* 2006;**107**:2S.

Sanchez-Ramos L, Gaudier FL, Kaunitz AM. Cervical ripening and labor induction after previous cesarean delivery. *Clinical Obstetrics and Gynecology* 2000;**43**:513-23.

Schmitz T, Goffinet F. [Against routine use of misoprostol in term labor induction]. *Gynecol.Obstet.Fertil.* 2006;**34**:161-5.